

Grail

The Bean Bag

A newsletter to promote communication among research scientists concerned with the systematics of the Leguminosae/Fabaceae

Number 49

February 2002

From the Editor

Barbara Mackinder

The Bean Bag is designed to promote communication among research scientists concerned with legume systematics. To achieve this goal *The Bean Bag* is issued annually. As of this volume, *The Bean Bag* will be issued in February each year. The change of month (from November) will allow for a more appropriate timing of certain regular announcements (e.g. The Rupert Barneby Award) and will avoid postal difficulties commonly associated with the Christmas period. *The Bean Bag* features six columns: From the Editor, News (meetings, major events, announcements, etc.), Latin American Legume Report, Nodulation and Nitrogen Fixation (new nodulation records), Gleanings, and Recent Legume Literature. Data in the Gleanings column are derived from questionnaire sheets which Readers complete and return.

The Recent Legume Literature column contains published research papers of specific interest to *Bean Bag* Readers and is derived from Readers contributions in conjunction with references from The Kew Record of Taxonomic Literature. Recent is defined as up to 18 months old. Specific interest to *Bean Bag* Readers is defined as research papers of interest to a worldwide group of legume systematic botanists.

Bean Bag Readers are encouraged to send notices, observations, etc. In particular in this issue, information is solicited concerning any aspect of nodulation (see the contribution from Janet Sprent under News). Also included is an inventory of legume taxa as yet unsampled for molecular phylogenetic studies (see the contribution from Gwilym Lewis and Brian Schrire under News). Furthermore a list of legume genera whose response to rhizobia is unknown is included with the intention of encouraging their exploration (see the contribution from Harold Corby and Joe Kirkbride under Nodulation and Nitrogen Fixation).

We encourage readers to accept delivery of *The Bean Bag* via e-mail to reduce our ever increasing distribution costs thus ensuring the continuation of *The Bean Bag*. If you are able to accept your copies by e-mail, please send a message to the editor (email: b.mackinder@rbgkew.org.uk). Will new readers please provide their title, first and last names, full postal address and area(s) of interest.

Electronic copies of the current and past issues of *Bean Bag* and the Current Readers Directory can be viewed on the World Wide Web server of the Royal Botanic Gardens, Kew, UK at <http://www.rbgkew.org.uk/herbarium/legumes/beanbag.html>

Bean Bag address:

Mrs B. Mackinder, Bean Bag Editor, Herbarium, Royal Botanic Gardens, Kew,
Richmond, Surrey TW9 3AB, United Kingdom
email: b.mackinder@rbgkew.org.uk

NEWS

The Fourth International Legume Conference, July 2001, Canberra, Australia

Joe Miller

The Fourth International Legume Conference, "Legumes Down Under", was held in Canberra during the first week of July 2001. There were over 100 participants at the conference with people from all over the world. The organisers were very pleased that there were very few logistical or technical problems and that the weather was cooperative. It was very useful time for legume workers to get together and to continue the work on legume phylogeny and other topics.

The conference had a strong focus on Systematics with other sessions on Phytochemistry, Utilization, Electronic and Web Based Tools, Symbiosis and Rehabilitation, Developmental and Structural Morphology and Legume/Animal Interactions.

Currently manuscripts, taken from papers given at the conference, are being reviewed for two publications of the Advances in Legume Systematics series. One volume (Advances X) will be published at The Royal Botanic Gardens, Kew and focuses on higher-level systematic studies. The second volume (Advances XI) will be published by CSIRO Publishing, will also be labeled as a special issue of Australian Systematic Botany and focuses on *Acacia* systematics and utilization.

Over a dozen participants also attended a post-conference symposium in Western Australia entitled "The Conservation and Utilization Potential of Dryland Acacias." A volume of papers from this symposium is also being prepared.

Several collaborative projects were discussed involving the Bean Tree Workshop and ILDIS. It is hoped that the conference has facilitated and reinforced cooperation and transfer of knowledge among legume workers and that this work will continue.

On behalf of the conference organizing committee Joe Miller would like to thank all the participants for visiting Canberra. We would also like to thank our many sponsors of the conference. We look forward to the next International Legume Conference and associated legume symposia at various meetings to continue our important work in legume research.

Organizing Committee: Joe Miller, Mike Crisp, Jim Grimes and David Morrison.

Nodulation in Legumes - not the end of the story!

Janet I. Sprent

Book Announcement

Janet Sprent's new book summarises the processes leading to nodulation in legumes and also the current list of those species known to nodulate. The last entry was made in March 2001, but there is already new information available. Two new genera of bacteria capable of nodulating legumes have been reported as well as another slant on the infection process. Much of the new information emerging is from the tropics and this is likely to be only the tip of the iceberg!

Please let Janet (jisprent@aol.com) know of any new material that you have, on any aspect of nodulation, so that it can be incorporated in an up-date of the book which is already planned.

Nodulation in Legumes by Janet I. Sprent (xii + 146pp. Soft cover, ISBN 1 84246 013 7) is published by The Royal Botanic Gardens, Kew, price £27.00 / €43.20 and can be purchased from <http://www.Kewbooks.com> and in the near future (expected April 2002) directly from RBG, Kew at <http://www.kew.org/publications>.

Nitrogen Fixation Conference

If you are working on nodulation in a developing country and would like to be considered for funding to attend the next European Conference on Nitrogen Fixation, to be held in Norwich, England Sept 6-10 2002, please email jisprent@aol.com with an indication of what you are doing and a brief CV.

A limited amount of funding has been obtained from the EU and it will be distributed competitively with a preference for young (post MSc or PhD) researchers.

The Leguminosae of Madagascar

Barbara Mackinder

The Leguminosae of Madagascar, edited by David J. Du Puy is the product of a project that originally aimed to produce a checklist of the Leguminosae of Madagascar. However it quickly became clear that the knowledge of the family in Madagascar was very incomplete and that a full flora account and fieldwork would be needed. Jean-Noel Labat, Jean Bosser and Jean-François Villiers from Paris collaborated on the book, as did Raymond Rabevohitra from Madagascar. Sadly J.-F. Villiers died before completion of the book. The book deals with 573 native species of which 80% are endemic to Madagascar. Six genera and 129 species were described as new to science during the project. The book is generously illustrated with line drawings and colour photographs. The Leguminosae of Madagascar, (720 pp. Hardcover, ISBN 1 900347 70 9) is published by The Royal Botanic Gardens, Kew. It can be purchased from <http://www.Kewbooks.com> and in the near future (expected April 2002) directly from RBG, Kew at <http://www.kew.org/publications> for £65.00 / €104.00 plus post/packaging.

Molecular sampling needs in Leguminosae

Gwilym P. Lewis and Brian D. Schrire

At a recent UK Legume group meeting, it was agreed that we should compile a list of genera that have not yet been sampled molecularly. In consultation with Marty Wojciechowski, Matt Lavin, Toby Pennington, Melissa Luckow and Pat Herendeen, we present below a list of critical genera (Table 1) for which no known molecular data is currently available.

If you are in a position to collect silica-dried leaf material of any of these genera (i.e. you have in place collecting permits for those areas where these genera are known to grow), please let Gwil Lewis or Brian Schrire at RBG, Kew know and they can put you in touch with the relevant specialist seeking to access this material. Alternatively, if you have already extracted DNA from any of these genera kindly also let us know so we can keep the list below updated. The overall aim of targeting these critical genera is to have available at least one species of each of the 720 genera of legumes, thus moving closer towards a full molecular legume phylogeny. We envisage this as Phase 1 of a legume gap-filling exercise, to be followed by a more detailed list including those paraphyletic or large genera where more detailed sampling is necessary. We present such a list for the Caesalpinioideae and tribe Mimoseae in Table 2 below.

Contact e-mails: G.Lewis@rbgkew.org.uk or B.Schrire@rbgkew.org.uk

Table 1. Genera for which no known molecular data is currently available

Genus	Tribe	Sp #	Distribution
CAESALPINIOIDEAE			
Arcoa	Caesalpineae	1	Haiti (Santo Domingo)
Balsamocarpon	Caesalpineae.	1	Chile
Chidlowia	Caesalpineae.	1	Sierra Leone, Ghana, Ivory Coast, Liberia
Moullava (=Wagatea)	Caesalpineae.	1	India
Neoapaloxylon	Caesalpineae.	1	Madagascar
Orphanodendron	Caesalpineae.	1	Colombia
Stenodrepanum	Caesalpineae.	1	Argentina
Stachyothyrsus	Caesalpineae.	3	Cameroon, Sierra Leone, Ivory Coast, Liberia, Zaire, Equatorial Guinea
Sympetalandra	Caesalpineae.	5	Indonesia & Malaysia (Sabah, Sarawak, Kalimantan, Malay Peninsula, Sumatra)
Adenolobus	Cercidieae	3	Namibia, S. Africa, Botswana
Gigasiphon	Cercidieae	3	Madagascar, Tanzania, Kenya, Angola, Gabon, Zaire
Griffonia	Cercidieae	4	Cameroon, Gabon, Nigeria, Zaire
Androcalymma	Cassieae	1	Brazil
Apuleia	Cassieae	1	Brazil, Colombia, Peru, Venezuela, Argentina
Eligmocarpus	Cassieae	1	Madagascar
Kalappia	Cassieae	1	Indonesia (Sulawesi) near Milili

Genus	Tribe	Sp #	Distribution
<i>Mendoravia</i>	Cassieae	1	Madagascar (southeast)
<i>Brandzeia</i> (= <i>Bathiaea</i>)	Detarieae s.l	1	Madagascar (southeast)
<i>Brachycylis</i>	Detarieae s.l	1	Colombia
<i>Brenaniodendron</i>	Detarieae s.l	1	Mozambique
<i>Brodriguesia</i>	Detarieae s.l	1	Brazil (Bahia)
<i>Hardwickia</i>	Detarieae s.l	1	west India
<i>Heterostemon</i>	Detarieae s.l	7	Guyana, Brazil (Amazonas, Para), Surinam, Colombia, Venezuela
<i>Lebruniodendron</i>	Detarieae s.l	1	Cameroon, Zaire
<i>Leucostegane</i>	Detarieae s.l	2	Malaysia & Indonesia (Malay Peninsula, Sumatra, Sarawak)
<i>Michelsonia</i>	Detarieae s.l	1	Zaire
<i>Neoapaloxylon</i>	Detarieae s.l	3	Madagascar (west and south)
<i>Pseudomacrolobium</i>	Detarieae s.l	1	Zaire
<i>Paloveopsis</i>	Detarieae s.l	1	Guyana, Brazil (Amazonas)
MIMOSOIDEAE			
<i>Aubrevillea</i>	Mimoseae	2	Guineo-Congolian
<i>Indopiptadenia</i>	Mimoseae	1	India & Nepal
<i>Lemurodendron</i>	Mimoseae	1	Madagascar
<i>Mimozgyanthus</i>	Mimoseae	1	Argentina
<i>Prosopidastrum</i>	Mimoseae	2	Baja, California & Argentina
<i>Xerocladia</i>	Mimoseae	1	South Africa, Namibia
<i>Stryphnodendron</i>	Mimoseae	c.30	Central America, northern South America and Brazil
<i>Abarema</i>	Ingeae	44	Mainly New World tropics, 1 sp. extending to Bahamas, 2 to SE Brazil
<i>Archidendron</i>	Ingeae	94	Sri Lanka, continental SE Asia, Malesia, NE Australia, 1 sp extending to Micronesia, 2 to Solomon Islands
<i>Archidendropsis</i>	Ingeae	14	New Guinea, Bismark Archipelago, Australia (Queensland), New Caledonia, Solomon Islands
<i>Blanchetiodendron</i>	Ingeae	1	E Brazil
<i>Cathormion</i>	Ingeae	1	SE Asia & Australia
<i>Cojoba</i>	Ingeae	12	West Indies, NW Andean and tras-Andean South America, SE Mexico & Central America
<i>Ebenopsis</i>	Ingeae	3	Mexico & S Texas
<i>Falcataria</i>	Ingeae	3	Moluccas, New Guinea, Solomon Islands, Australia (Queensland)
<i>Guinetia</i>	Ingeae	1	Mexico
<i>Havardia</i>	Ingeae	5	Mexico, Texas & C America
<i>Hesperalbizia</i>	Ingeae	1	SW Mexico
<i>Hydrochorea</i>	Ingeae	3-4	Orinoco & Amazon basins, Guianas, Brazil
<i>Leucochloron</i>	Ingeae	4	Brazil
<i>Macrosamanea</i>	Ingeae	11	South America, mostly Amazonian
<i>Painteria</i>	Ingeae	3	Mexico
<i>Serianthes</i>	Ingeae	c.18	Thailand, Malesia, Micronesia, Melanesia & W. Polynesia
<i>Sphinga</i>	Ingeae	3	Mexico, Cuba, Guatemala, & NW South America
<i>Viguieranthus</i>	Ingeae	23	Asia & Madagascar
<i>Wallaceodendron</i>	Ingeae	1	N Celebes & Philippines
<i>Zygia</i>	Ingeae	c.60	Lowland tropical America and Greater Antilles to South America, most diverse in Central America, Columbia, Guianas, & NW Amazonia
PAPILIONOIDEAE			Genera arranged systematically within tribes
<i>Candolleodendron</i>	Swartzieae	1	S America (French Guiana, N and NE Brazil)
<i>Haplormosia</i>	Sophoreae	1	W and C Africa (Sierra Leone to Gabon)
<i>Uleanthus</i>	Sophoreae	1	S America (Amazonian Brazil)
<i>Panurea</i>	Sophoreae	2	S America (Colombia and Brazil)

Helen
1
Helen
3

See e-mail of May 24

Genus	Tribe	Sp #	Distribution
<i>Diplostropis</i>	Sophoreae	12	S America, (Colombia and Venezuela to Peru and Bolivia)
<i>Spirotropis</i> <i>Helen I.</i>	Sophoreae	2-3	Northern S America (Venezuela, Guyana, Surinam & French Guiana)
<i>Monopteryx</i>	Sophoreae	3-4	Northern S America (Colombia, Venezuela, French Guiana and Amazonian Brazil)
<i>Petaladenium</i>	Sophoreae	1	Brazil (Rio Negro river)
<i>Sakoanala</i>	Sophoreae	2	Madagascar
<i>Neoharussia</i> <i>Helen I.</i>	Sophoreae	2	N and NW Madagascar
<i>Ammothamnus</i>	Sophoreae	3	Central Asia (Kazakstan, Tadzhikistan, Uzbekistan and Turkmenistan)
<i>Amphiurus</i>	Sophoreae	3-4	C and W Africa
<i>Baphiastrum</i>	Sophoreae	1	Tropical central Africa
<i>Ammopiptanthus</i>	Thermop-sidae	1-2	Mongolia. China, Russia
<i>Xiphotheca</i>	Podalyriae	9	Africa (endemic to the Cape region of South Africa)
<i>Amphithalea</i>	Podalyriae	42	Africa (endemic to the Cape region of South Africa)
<i>Pearsonia</i>	Crotalariaeae	13	Africa and Madagascar
<i>Rothia</i>	Crotalariaeae	2	Africa and Asia to Australia
<i>Robynsiophyton</i>	Crotalariaeae	1	SC Africa (Angola, Zambia and Zaire)
<i>Spartidium</i>	Crotalariaeae	1	N Africa (Morocco to Libya)
<i>Bolusia</i>	Crotalariaeae	5	SC and southern Africa
<i>Wiborgia</i>	Crotalariaeae	10	Africa (endemic to the Cape region of South Africa)
<i>Polhillia</i>	Genisteae	7	W and SW parts of the Cape region of South Africa
<i>Sellocharis</i>	Genisteae	1	S America (SE Brazil)
<i>Anarthrophyllum</i>	Genisteae	15	S America (Andes in Argentina and Chile)
<i>Hesperolaburnum</i>	Genisteae	1	Morocco (Anti Atlas Mts.)
<i>?Cytisophyllum</i>	Genisteae	1	Mediterranean area of S Europe from Spain to Italy
<i>Cytisopsis</i>	Loteae	1-2	Mediterranean (E Mediterranean and N Africa)
<i>Antopetitia</i>	Loteae	1	Tropical Africa (mountains)
<i>Oreophysa</i>	Galegeae	1	N. Iran, mountains
<i>Ereinosparton</i>	Galegeae	3	SE Russia and Caucasus to C Asia and NW China
<i>Calophaca</i>	Galegeae	5	Central Asia, Ukraine to Caucasus
<i>Chesneya</i>	Galegeae	c.30	Temperate SW Asia to Sino-Himalayan region, most diverse in C Asia to Mongolia
<i>Neodielsia</i> (<i>Astragalus</i> segr.)	Galegeae	?	Turkestan
<i>Smirnowia</i>	Galegeae	1	Afganistan, Iran, Turkestan
<i>Gueldenstaedtia</i>	Galegeae	c. 10	Sino-Himalayan region to Mongolia and Siberia
<i>Stracheya</i>	Hedysareae	1	High Himalayas (Kashmir, Tibet, etc.)
<i>Taverniera</i>	Hedysareae	10	W India to Middle East, Horn of Africa
<i>Sartoria</i>	Hedysareae	1	S Turkey
<i>Ebenus</i>	Hedysareae	20	Mediterranean
<i>Eversmannia</i>	Hedysareae	1	Iran to W China
<i>Vavilova</i>	Fabeae	1	Turkey, Middle East
<i>Parochetus</i>	Trifolieae	2	Mtns. of tropical East Africa and Himalayas (no sampling from natural collections)
<i>Endosamara</i>	Millettieae	1	Indian subcontinent, Indo-China and Malasia
<i>Sarcodum</i>	Millettieae	c.3	S. China, Indo-China, Malesia, Papuasias
<i>Burkilliodendron</i>	Millettieae	1	Malaya (Perak)
<i>Craspedolobium</i>	Millettieae	1	S and SW China, Indo-China (Burma, Thailand, Laos)
<i>Kunstleria</i>	Millettieae	8	Malesia and Indian subcontinent (1 sp.)
<i>Schefflerodendron</i>	Millettieae	4	West-Central Tropical Africa with 1 sp. to East Africa
<i>Disynstemon</i>	Millettieae	1	Madagascar (South West)
<i>Pyranthus</i>	Millettieae	6	W, S and C Madagascar
<i>Sylvichadsia</i>	Millettieae	4	E and N Madagascar

Genus	Tribe	Sp #	Distribution
<i>Paratephrosia</i>	Millettieae	1	Australia (W. Australia, N. Territory, S. Australia, Queensland)
<i>Ptycholobium</i>	Millettieae	3	W to NE Tropical Africa and Arabia; southern Africa
<i>Requienia</i>	Millettieae	3	W to NE Tropical Africa (Sahelian zone); southern Africa
<i>Platysepalum</i>	Millettieae	7-8	Tropical, mostly West Africa
<i>Antheroporum</i>	Millettieae	c.3	SW China and Indo-China (Burma, Thailand and Vietnam)
<i>?Imbralyx</i>	Millettieae	c.4	S China to Sumatra (segregate of <i>Fordia</i>)
<i>Behaimia</i>	Millettieae	1	Cuba
<i>Bergeronia</i>	Millettieae	1	Brazil, Bolivia, Paraguay and Argentina
<i>Dahlstedtia</i>	Millettieae	2	Brazil, Argentina
<i>Margaritolobium</i>	Millettieae	1	Venezuela
<i>Meizotropis</i>	Phaseoleae	2	tropical Asia (subcontinental India and W Indo-China)
<i>Cochlianthus</i>	Phaseoleae	2	W China (Yunnan, Sichuan) and Himalayas (Nepal)
<i>Cruddasia</i>	Phaseoleae	c.2	NE Indian subcontinent; Indo-China (Burma, Thailand)
<i>Pseudoeriosema</i>	Phaseoleae	c.4	tropical Africa
<i>Barbieria</i>	Phaseoleae	1	Neotropics, from S Mexico, C America, Caribbean and W tropical S America
<i>Clitoriopsis</i>	Phaseoleae	1	tropical Africa (Congo [Kinshasa] and Sudan)
<i>Macropsychanthus</i>	Phaseoleae	c.2	Papuasias, Micronesia (possibly Philippines)
<i>?Luzonia</i>	Phaseoleae	1	Philippines (Luzon and Leyte)
<i>Herpyza</i>	Phaseoleae	1	Cuba (Allen & Allen, 1981 also note its occurrence in Puerto Rico)
<i>Diphyllarium</i>	Phaseoleae	1	Indo-China (Laos & Vietnam)
<i>Sinodolichos</i>	Phaseoleae	2	SE Asia (NE Indian subcontinent, SW China, Indo-China)
<i>Nogra</i>	Phaseoleae	c.3	Indian subcontinent, S China, Indo-China (all 3 spp. should be sampled)
<i>Dysolobium</i>	Phaseoleae	4	SE Asia (E Indian subcontinent, Indo-China, SW China, Malesia)
<i>Alistilis</i>	Phaseoleae	3	Southern Africa (1 sp.), Madagascar (2 spp.)
<i>Nesphostylis</i>	Phaseoleae	4	tropical and subtropical Africa (2 spp.); SE Asia (Indian subcontinent, Indo-China [Burma]; 2 spp.)
<i>Decorsea</i>	Phaseoleae	6	Southern tropical and subtropical Africa (3 spp.) and Madagascar (3 spp.)
<i>Spathionema</i>	Phaseoleae	1	E tropical Africa
<i>Vatovaea</i>	Phaseoleae	1	NE tropical Africa
<i>Dunbaria</i>	Phaseoleae	20	SE Asia (Indian subcontinent, Indo-China, China, E Asia, Malesia, Papuasias); N Australia
<i>Bolusafr</i>	Phaseoleae	1	South Africa (W Cape)
<i>Flemingia</i>	Phaseoleae	30-35	SE Asia (Indian subcontinent, Indo-China, China, Malesia, Papuasias) to Australia (ca 28-33 spp.); Africa 1 sp. and <i>F. grahamiana</i> Wight and Arn. widespread in Africa and Asia
<i>Paracalyx</i>	Phaseoleae	6	NE tropical Africa (Ethiopia, Somalia) and Socotra (5 spp.); Indian subcontinent, Indo-China (1 sp.)
<i>Carissoa</i>	Phaseoleae	1	SW tropical Africa (Angola)
<i>Chrysocias</i>	Phaseoleae	3-4	South Africa (S parts of W Cape)
<i>Orbexilum</i>	Psoraleae	8	E USA from Virginia to Florida, westwards through Kansas, Oklahoma and Texas; Mexico (to Chiapas)
<i>Hoita</i>	Psoraleae	3	W USA (California) to NW Mexico (Baja California)
<i>Pediomelum</i>	Psoraleae	21	SC Canada, USA (Great Plains and Great Basin) to C Mexico

Table 2: Genera for which additional species are needed

CAESALPINIOIDEAE			
Genus	Tribe	Sp #	Distribution
<i>Caesalpinia s.l.</i>	Caesal.	c.100	pantropical
<i>Mezoneuron</i>	Caesal.	c.30	Malaysia & Indonesia (Malay Peninsula, Java, Sumatra, Sarawak, Kalimantan, Celebes), Thailand, trop Africa, Hawaii
<i>Tachigali s. l.</i>	Caesal.	60	C & S. America
<i>Dimorphandra</i>	Caesal.	25	Guyana, Surinam, Brazil (Amazonas, Para, Rio de Janeiro), Venezuela, Colombia
<i>Gleditsia</i>	Caesal.	14	Indonesia (Celebes)
<i>Bauhinia</i>	Cercid.	c.250	pantropical
<i>Cassia</i>	Cassieae	30	pantropical
<i>Dialium</i>	Cassieae	c.40	pantropical
<i>Labichea</i>	Cassieae	18	Australia (Queensland, W. Austr.)
<i>Martiodendron</i>	Cassieae	4	Brazil (Para, Amazonas, Bahia, Maranhao), Guyana
<i>Senna</i>	Cassieae	240	pantropical
<i>Brachystegia</i>	Detarieae s.l	c.30	west & east Africa
<i>Brownea</i>	Detarieae s.l	c.30	Guyana, French Guiana, Venezuela, Ecuador, Colombia, Peru
<i>Crudia</i>	Detarieae s.l	c.55	Malaysia & Indonesia (Sumatra, Java, Sabah, Kalimantan, Sarawak, Malay Peninsula), Guyana, Brazil (Para, Amazonas, Rondonia), Fr. Guiana, Surinam, Colombia, Zaire
<i>Daniellia</i>	Detarieae s.l	9	Tanzania, west Africa including Cameroon, Gabon, Zaire
<i>Dicymbe</i>	Detarieae s.l	15	Guyana, Brazil, Peru, Venezuela
<i>Elizabetha</i>	Detarieae s.l	10	Guyana, Surinam, French Guiana, Brazil
<i>Guibourtia</i>	Detarieae s.l	c.17	Brazil, Paraguay, Caribbean, Sierra Leone, Gabon, Congo, Zaire, Zambia, Mozambique, S. Africa
<i>Macrolobium</i>	Detarieae s.l	c.100	Guyana, Fr. Guiana, Surinam, Ecuador, Brazil, Peru, Venezuela, Colombia
<i>Maniltoa</i>	Detarieae s.l	c.25	Malaysia & Indonesia (Malay Peninsula, Celebes), New Guinea, , Australia
<i>Paloue</i>	Detarieae s.l	4	Guyana, French Guiana, Surinam
<i>Peltogyne</i>	Detarieae s.l	23	Guyana, Fr. Guiana, Surinam, Brazil, Peru, Mexico, W. Indies
<i>Sindora</i>	Detarieae s.l	18-20	Malaysia & Indonesia (Malay Peninsula, Sabah, Kalimantan, Sarawak, Sumatra, Celebes, Java
<i>Zenkerella</i>	Detarieae s.l	5	Tanzania, Cameroon, Gabon
MIMOSOIDEAE			
Genus	Tribe	Sp #	Distribution
<i>Piptadenia</i>	Mimoseae	c.21	Mexico, throughout Central and South America as far south as Argentina and peru
<i>Prosopis</i> (especially <i>P. africana</i>)	Mimoseae	c.44	SW North America, South America, SW Asia & Africa
<i>Mimosa</i> (especially sect. <i>Mimadenia</i>)	Mimoseae	c.480	mostly New World tropics, a few in tropical Africa, S Asia and Madagascar

The Rupert Barneby Award

James L. Luteyn

The New York Botanical Garden is pleased to announce that Aaron Liston, currently at the Department of Botany & Plant Pathology, Oregon State University, is the recipient of the Rupert Barneby Award for the year 2002. Dr. Liston will be studying the phylogenetic systematics of *Astragalus* and *Trifolium*.

The New York Botanical Garden now invites applications for the Rupert Barneby Award for the year 2003. The award of US\$ 1,000.00 is to assist researchers to visit The New York Botanical Garden to study the rich collection of Leguminosae. Anyone interested in applying for the award should submit their curriculum vitae, a detailed letter describing the project for which the award is sought, and the names of 2-3 referees. Travel to the NYBG should be planned for sometime in the year 2003. The application should be addressed to Dr. James L. Luteyn, Institute of Systematic Botany, The New York Botanical Garden, 200th Street and Kazimiroff Blvd., Bronx, NY 10458-5126 USA, and received no later than December 1, 2002. Announcement of the recipient will be made by December 15th.

Anyone interested in making a contribution to THE RUPERT BARNEBY FUND IN LEGUME SYSTEMATICS, which supports this award, may send their check, payable to The New York Botanical Garden, to Dr. Luteyn.

Rupert Barneby, 1911–2000 (Obituary)

Gwilym P. Lewis

Rupert was born on October 6, 1911, in Monmouthshire, on the English-Welsh border. He went to harrow Public School from 1924–1929 and to Trinity College, Cambridge University from 1930–1932 where he gained a B. A. in History and Modern Languages.

One of Rupert's early enthusiasms in legumes when collecting in Spain and N. Africa prior to the Hitler war was the *Cytisus-Ulex-Stauracanthus* complex and it is evident that he entered the United States in 1937 with beans already a main focus of his botanical interest. He took up permanent residency in the United States in 1941.

His main research interests were xerophytic Floras, the taxonomy of the Leguminosae and Neotropical Menispermaceae. He described 1,160 species new to science in upwards of 140 publications, all of which are learned papers and many of which are monumental tomes. Of the 25 species named by other botanists in his honour, over a third are species of legume. Four genera honour him, and three of these are legumes, *Barnebydendron* Kirkbride, *Barnebyella* Podlech and *Rupertia* Grimes. Eight of his own legume publications (some coauthored): North American *Astragalus*, Daleae Images, The American Cassiinae, A monograph of *Mimosa*, A conspectus of *Erythrina* and three volumes on the New World Ingeae (including his last major work, *Calliandra*) describe in total 1,900 species.

Rupert's passing leaves an echoing gap in New York and in the world of botany.

Bean Bag readers' attention is drawn to "Ruperti Images: a portrait of Rupert Barneby" by Douglas Crase in *Brittonia* 53(1): 1–40 (2001).

Nodulation and Nitrogen Fixation

(Legume Nodulation reports not in Allen and Allen (1981))

Joseph H. Kirkbride, Jr.

Taxon	Status ¹	Source ²
<i>Abarema bigemina</i> (L.) Kosterm.	+	164
<i>Abarema cochleata</i> (Willd.) Barneby & J.W. Grimes var. <i>moniliformis</i> (Ducke) Barneby & J.W. Grimes	+	38
<i>Abarema cochlicarpos</i> (B.A. Gomes) Barneby & J.W. Grimes	+	50
<i>Abrus canescens</i> Baker	+	49
<i>Abrus cantoniensis</i> Hance	+	72
<i>Abrus mollis</i> Hance	+	72
<i>Abrus pulchellus</i> Wall. ex Thwaites subsp. <i>tenuiflorus</i> (Benth.) Verdc.	+	42
<i>Acacia acradenia</i> F. Muell.	+	17
<i>Acacia amoena</i> H.L. Wendl.	+	138
<i>Acacia ancistrocarpa</i> Maiden & Blakely	+	17
<i>Acacia argyrodendron</i> Domin	+	20
<i>Acacia atramentaria</i> Benth.	+	124
<i>Acacia berlandieri</i> Benth.	+	69
<i>Acacia binervata</i> DC.	+	170
<i>Acacia bivenosa</i> DC.	+	45
<i>Acacia blakei</i> Pedley	+	138
<i>Acacia brassii</i> Pedley	+	36
<i>Acacia brevispica</i> Harms	+	20
<i>Acacia brumalis</i> Maslin	+	35
<i>Acacia burrowii</i> Maiden	+	138
<i>Acacia caesia</i> (L.) Willd.	+	150
<i>Acacia cambagei</i> R.T. Baker	+	138
<i>Acacia cangaiensis</i> Tindale & Kodela	+	100
<i>Acacia chrysotricha</i> Tindale	+	170
<i>Acacia cincinnat</i> F. Muell.	+	36
<i>Acacia citrinoviridis</i> Tindale & Maslin	+	150
<i>Acacia colei</i> Maslin & A.J. Thomson	+	144
<i>Acacia concinna</i> (Willd.) DC.	+	57
<i>Acacia concinna</i> (Willd.) DC.	-	76
<i>Acacia concurrens</i> Pedley	+	38
<i>Acacia coriacea</i> DC.	+	20
<i>Acacia cowleana</i> Tate	+	17
<i>Acacia cuthbertsonii</i> Luehm.	+	17
<i>Acacia cyperophylla</i> F. Muell. ex Benth.	+	20
<i>Acacia dangarensis</i> Tindale & Kodela	+	25
<i>Acacia dictyophleba</i> F. Muell.	+	17
<i>Acacia difficilis</i> Maiden	+	144
<i>Acacia dimidiata</i> Benth.	+	138
<i>Acacia dolichostachya</i> S.F. Blake	+	150
<i>Acacia drepanolobium</i> Harms ex Y. Sjostedt	+	13
<i>Acacia dudgeoni</i> Holland	+	74
<i>Acacia dunnii</i> (Maiden) Turrill	+	138
<i>Acacia elatior</i> Brenan	+	121
<i>Acacia elongata</i> Sieber ex DC.	+	138
<i>Acacia eremaea</i> C.R.P. Andrews	+	35
<i>Acacia erioloba</i> E. May.	+	64
<i>Acacia etbaica</i> Schweinf.	+	7

Ed. Note: Reference 150 is "Nodulation in Legumes" by Janet I. Sprent published by the Royal Botanic Gardens, Kew (2001).

Taxon	Status ¹	Source ²
<i>Acacia falciformis</i> DC.	+	170
<i>Acacia ferruginea</i> DC.	+	31
<i>Acacia filicifolia</i> Cheel & M.B. Welch	+	138
<i>Acacia flavescens</i> Benth.	+	20
<i>Acacia fulva</i> Tindale	+	170
<i>Acacia gittinsii</i> Pedley	+	138
<i>Acacia glauca</i> Moench	+	72
<i>Acacia glaucocarpa</i> Maiden & Blakely	+	100
<i>Acacia gonocarpa</i> F. Muell.	+	21
<i>Acacia gourmaensis</i> A. Chev.	+	74
<i>Acacia gummiifera</i> Willd.	+	94
<i>Acacia hemsleyi</i> Maiden	+	17
<i>Acacia hillianiana</i> Maiden	+	150
<i>Acacia hippuroides</i> Heward ex Benth.	+	150
<i>Acacia hockii</i> De Wild.	+	14
<i>Acacia holosericea</i> A. Cunn. ex G. Don	+	69
<i>Acacia holosericea</i> A. Cunn. ex G. Don var. <i>neurocarpa</i> (Hook.) Domin	+	144
<i>Acacia implexa</i> Benth.	+	138
<i>Acacia irrorata</i> Sieber ex Spreng.	+	170
<i>Acacia irrorata</i> Sieber ex Spreng. ssp. <i>irrorata</i>	+	170
<i>Acacia irrorata</i> Sieber ex Spreng. ssp. <i>velutinella</i> Tindale	+	170
<i>Acacia iteaphylla</i> F. Muell. ex Benth.	+	138
<i>Acacia ixiophylla</i> Benth.	+	35
<i>Acacia julifera</i> Benth.	+	36
<i>Acacia lachnophylla</i> F. Muell.	+	138
<i>Acacia laeta</i> R. Br. ex Benth.	+	74
<i>Acacia latescens</i> Benth.	+	104
<i>Acacia leiocalyx</i> (Domin) Pedley	+	138
<i>Acacia leucoclada</i> Tindale	+	21
<i>Acacia linarioides</i> Benth.	+	45
<i>Acacia litakumensis</i> Burch.	+	20
<i>Acacia longispicata</i> Benth.	+	144
<i>Acacia lysiphloia</i> F. Muell.	+	17
<i>Acacia maconochieana</i> Pedley	+	36
<i>Acacia melleodora</i> Pedley	+	17
<i>Acacia minuta</i> (M.E. Jones) R.M. Beauch.	+	6
<i>Acacia montigena</i> Brenan & Exell	+	150
<i>Acacia mountfordiae</i> Specht	+	138
<i>Acacia mutabilis</i> Maslin ssp. <i>mutabilis</i>	+	35
<i>Acacia mucronata</i> Willd. ex H.L. Wendl.	+	104
<i>Acacia mutabilis</i> Maslin ssp. <i>stipulifera</i> Maslin	+	35
<i>Acacia nanodealbata</i> J.H. Willis	+	170
<i>Acacia negrii</i> Pic. Serm.	+	7
<i>Acacia neurocarpa</i> Hook.	+	144
<i>Acacia nilotica</i> (L.) Willd. ex Delile ssp. <i>tomentosa</i> (Benth.) Brenan	+	45
<i>Acacia notabilis</i> F. Muell.	+	138
<i>Acacia oncinocarpa</i> Benth.	+	138
<i>Acacia oshanesii</i> F. Muell. & Maiden	+	63
<i>Acacia papyrocarpa</i> Benth.	+	138
<i>Acacia parvipinnula</i> Tindale	+	170
<i>Acacia patagiata</i> R.S. Cowan & Maslin	+	35
<i>Acacia pellita</i> O. Schwarz	+	105
<i>Acacia pendula</i> A. Cunn. ex G. Don	+	138
<i>Acacia pentagona</i> (Schumacher & Thonn.) Hook. f.	+	68
<i>Acacia planifrons</i> Wight & Arn.	+	31
<i>Acacia plectocarpa</i> A. Cunn.	+	36
<i>Acacia polyacantha</i> Willd. ssp. <i>polyacantha</i>	+	31
<i>Acacia polystachya</i> Benth.	+	36

Taxon	Status ¹	Source ²
<i>Acacia prasinata</i> Asfaw	+	7
<i>Acacia pyrifolia</i> DC.	+	138
<i>Acacia redolens</i> Maslin	+	35
<i>Acacia reficiens</i> Wawra	+	121
<i>Acacia rothii</i> F.M. Bailey	+	36
<i>Acacia rosumae</i> Oliv.	+	74
<i>Acacia sakalava</i> Drake	-	74
<i>Acacia schaffneri</i> (S. Watson) F.J. Herm.	+	150
<i>Acacia schottii</i> Torr.	+	150
<i>Acacia schweinfurthii</i> Brenan & Exell	-	68
<i>Acacia shirleyi</i> Maiden	+	20
<i>Acacia silvestris</i> Tindale	+	170
<i>Acacia simsii</i> Benth.	+	36
<i>Acacia sparsiflora</i> Maiden	+	20
<i>Acacia spectabilis</i> A. Cunn. ex Benth.	+	36
<i>Acacia stipuligera</i> F. Muell.	+	17
<i>Acacia sumia</i> (Roxb.) Buch.-Ham. ex J. Voigt	+	31
<i>Acacia sutherlandii</i> (F. Muell.) F. Muell.	+	20
<i>Acacia teniana</i> Harms	+	72
<i>Acacia tenuissima</i> F. Muell.	+	17
<i>Acacia tephrrina</i> Pedley	+	20
<i>Acacia thomsonii</i> Maslin & M.W. McDonald	+	17
<i>Acacia tortilis</i> (Forssk.) Hayne	+	69
<i>Acacia torulosa</i> Benth.	+	104
<i>Acacia trachycarpa</i> E. Pritz.	+	134
<i>Acacia trachyphloia</i> Tindale	+	21
<i>Acacia translucens</i> A. Cunn. ex Hook.	+	150
<i>Acacia trineura</i> F. Muell.	+	138
<i>Acacia tumida</i> F. Muell. ex Benth.	+	20
<i>Acacia ulicifolia</i> (Salisb.) Court var. <i>brownei</i> (Steud.) Pedley	+	159
<i>Acacia umbellata</i> Benth.	+	21
<i>Acacia verniciflua</i> A. Cunn.	+	138
<i>Acacia vestita</i> Ker Gawl.	+	138
<i>Acacia villosa</i> (Sw.) Willd.	+	20
<i>Acacia welwitschii</i> Oliv.	+	69
<i>Acacia yirrkallensis</i> Specht	+	104
<i>Acacia zanzibarica</i> (S. Moore) Taub.	+	121
<i>Acrocarpus fraxinifolius</i> Wight ex Arn.	+	69
<i>Acrocarpus fraxinifolius</i> Wight ex Arn.	-	50
<i>Adenanthera intermedia</i> Merr.	-	68
<i>Adenanthera microsperma</i> Teijsm. & Binn.	+	130
<i>Aeschynomene ciliata</i> Vogel	+	4
<i>Aeschynomene crassicaulis</i> Harms	+	4
<i>Aeschynomene cristata</i> Vatke	+	98
<i>Aeschynomene cristata</i> Vatke var. <i>pubescens</i> J. Léon	+	145
<i>Aeschynomene denticulata</i> Rudd	+	46
<i>Aeschynomene pfundii</i> Taub.	+	4
<i>Aeschynomene pratensis</i> Small	+	46
<i>Aeschynomene scabra</i> G. Don	+	46
<i>Aeschynomene tambacoundensis</i> Berhaut	+	4
<i>Aeschynomene uniflora</i> E. Mey.	+	42
<i>Aeschynomene villosa</i> Poir.	+	4
<i>Afzelia quanxensis</i> Welw.	+	78
<i>Albizia amara</i> (Roxb.) Boivin	+	69
<i>Albizia berteriana</i> (Balb. ex DC.) M. Gomez	+	69
<i>Albizia chevalieri</i> Harms	+	165
<i>Albizia dinklagei</i> (Harms) Harms	-	10
<i>Albizia edwallii</i> (Hoehne) Barneby & J.W. Grimes	+	50

Taxon	Status ¹	Source ²
<i>Albizia ferruginea</i> (Guill. & Perr.) Benth.	+	163
<i>Albizia glaberrima</i> (Schumach. & Thonn.) Benth.	+	69
<i>Albizia kalkora</i> Prain	+	72
<i>Albizia kalkora</i> Prain	-	108
<i>Albizia lucidor</i> (Steud.) I.C. Nielsen ex H. Hara	+	76
<i>Albizia petersiana</i> (Bolle) Oliv.	+	69
<i>Albizia schimperana</i> Oliv.	+	69
<i>Albizia suluensis</i> Gerstner	+	61
<i>Albizia zygia</i> (DC.) J.F. Macbr.	+	163
<i>Alhagi sparsifolia</i> (Shap.) Shap.	+	72
<i>Amicia zygotomeris</i> DC.	-	150
<i>Amphicarpaea trisperma</i> Miq.	+	155
<i>Amphimas pterocarpoides</i> Harms	-	150
<i>Amphithalea ericaefolia</i> (L.) Eckl. & Zeyh.	+	61
<i>Anadenanthera colubrina</i> (Vell.) Brenan	-	6
<i>Anadenanthera colubrina</i> (Vell.) Brenan var. <i>cebil</i> (Griseb.) Altschul	-	6
<i>Anadenanthera falcata</i> (Benth.) Speg.	+	87
<i>Anadenanthera peregrina</i> (L.) Speg. var. <i>falcata</i> (Benth.) Altschul	+	87
<i>Andira anthelmia</i> (Vell.) J.F. Macbr.	+	50
<i>Andira nitida</i> Mart. ex Benth.	+	53
<i>Andira spectabilis</i> Saldanha	+	68
<i>Angylocalyx oligophyllus</i> (Baker) Baker f.	-	150
<i>Anthonotha fragrans</i> (Baker f.) Exell & Hillc.	-	77
<i>Anthyllis cytisoides</i> L.	+	158
<i>Anthyllis vulneraria</i> L. ssp. <i>polyphylla</i> (DC.) Nyman	+	112
<i>Apios carnea</i> (Wall.) Benth. ex Baker	+	76
<i>Apuleia leiocarpa</i> (Vogel) J.F. Macbr.	+	68
<i>Arachis batizocoi</i> Krapov. & W.C. Greg.	+	166
<i>Arachis burkartii</i> Handro	+	28
<i>Arachis cardenasii</i> Krapov. & W.C. Greg.	+	166
<i>Arachis glabrata</i> Benth. var. <i>hagenbeckii</i> (Harms) F.J. Herm.	+	150
<i>Arachis helodes</i> Mart. ex Krapov. & W.C. Greg.	+	47
<i>Arachis monticola</i> Krapov. & Rigoni	+	166
<i>Arachis pintoii</i> Krapov. & W.C. Greg.	+	38
<i>Arachis pusilla</i> Benth.	+	28
<i>Arachis stenosperma</i> Krapov. & W.C. Greg.	+	90
<i>Archidendron basaltica</i> (F. Muell.) I.C. Nielsen	+	20
<i>Archidendron bigeminum</i> (L.) I.C. Nielsen	+	164
<i>Archidendron clypearia</i> (Jack) I.C. Nielsen	-	71
<i>Archidendron kanisii</i> R.S. Cowan	+	150
<i>Archidendropsis thozetiana</i> (F. Muell.) I.C. Nielsen	+	23
<i>Argyrolobium filiforme</i> Eckl. & Zeyh.	+	61
<i>Argyrolobium roseum</i> ssp. <i>ornithopodioides</i> (Jaub. & Spach.) Jafri & Ali	+	9
<i>Aspalathus acuminata</i> Lam. ssp. <i>pungens</i> (Thunb.) Dahlgren	+	61
<i>Aspalathus asparagoides</i> L. f. ssp. <i>asparagoides</i>	+	61
<i>Aspalathus chortophila</i> Eckl. & Zeyh. Ssp. <i>kougaensis</i> Dahlgren	+	61
<i>Aspalathus confusa</i> Dahlgren	+	61
<i>Aspalathus divaricata</i> Thunb.	+	40
<i>Aspalathus forbesii</i> Harv.	+	40
<i>Aspalathus frankenioides</i> DC.	+	61
<i>Aspalathus heterophylla</i> L. f. ssp. <i>heterophylla</i>	+	61
<i>Aspalathus hirta</i> E. Mey. ssp. <i>hirta</i>	+	61
<i>Aspalathus hispida</i> Thunb.	+	40
<i>Aspalathus lactea</i> Thunb. ssp. <i>breviloba</i> Dahlgren	+	61
<i>Aspalathus lactea</i> Thunb. ssp. <i>lactea</i>	+	61
<i>Aspalathus longifolia</i> Benth.	+	61
<i>Aspalathus longipes</i> Harv.	+	61
<i>Aspalathus nivea</i> Thunb.	+	60

Taxon	Status ¹	Source ²
<i>Aspalathus opaca</i> Eckl. & Zeyh. ssp. <i>rostriloba</i> Dahlgren	+	61
<i>Aspalathus pachyloba</i> Benth. ssp. <i>macroclada</i> Dahlgren	+	61
<i>Aspalathus quinquefolia</i> L. ssp. <i>virgata</i> (Thunb.) Dahlgren	+	61
<i>Aspalathus retroflexa</i> L. ssp. <i>bicolor</i> (Eckl. & Zeyh.) Dahlgren	+	61
<i>Aspalathus spicata</i> Thunb. ssp. <i>cliffortioides</i> (Schltr.) Dahlgren	+	61
<i>Aspalathus spinosa</i> L. ssp. <i>flavisipina</i> (Presl ex Benth.) Dahlgren	+	61
<i>Aspalathus spinosa</i> L. ssp. <i>glauca</i> (Eckl. & Zeyh.) Dahlgren	+	61
<i>Aspalathus spinosissima</i> R. Dahlgren	+	33
<i>Aspalathus ternata</i> (Thunb.) Druce	+	61
<i>Aspalathus tuberculata</i> Walp.	+	61
<i>Astragalus complanatus</i> R. Br. ex Bunge	+	162
<i>Astragalus coquimbensis</i> (Hook. & Arn.) Reiche	+	59
<i>Astragalus coronilloides</i> Ulbr.	+	162
<i>Astragalus danicus</i> Retz.	+	127
<i>Astragalus densiflorus</i> Kar. & Kir.	+	169
<i>Astragalus donianus</i> DC.	+	76
<i>Astragalus echinatus</i> Murray	+	150
<i>Astragalus epiglottis</i> L.	+	150
<i>Astragalus falciformis</i> Desf.	+	150
<i>Astragalus glaux</i> L.	+	150
<i>Astragalus gombo</i> Bunge	+	150
<i>Astragalus graveolens</i> Benth.	+	117
<i>Astragalus himalayanus</i> Klotzsch	+	150
<i>Astragalus jaegerianus</i> Munz	+	59
<i>Astragalus luristanicus</i> Freyn	+	146
<i>Astragalus miniatus</i> Bunge	+	162
<i>Astragalus palmeri</i> A. Gray	+	48
<i>Astragalus persepolitanus</i> Boiss.	+	9
<i>Astragalus politovii</i> Krylov	+	120
<i>Astragalus scaberrimus</i> Bunge	+	162
<i>Astragalus sesameus</i> L.	+	150
<i>Astragalus stevenianus</i> DC.	+	162
<i>Astragalus tibetanus</i> Benth. ex Bunge	+	76
<i>Astragalus tribuloides</i> Delile	+	9
<i>Astragalus uliginosus</i> L.	+	120
<i>Astragalus xiphocarpus</i> Benth. ex Bunge	+	76
<i>Ateleia apetala</i> Griseb.	+	150
<i>Ateleia gummiifera</i> (DC.) D. Dietr.	+	150
<i>Baphia bequaertii</i> De Wild.	+	77
<i>Baphia laurifolia</i> Baill.	+	77
<i>Baphia maxima</i> Baker	+	150
<i>Baphia pubescens</i> Hook. f.	+	122
<i>Baphiopsis parviflora</i> Benth. ex Baker	-	150
<i>Bauhinia blakeana</i> Dunn	-	69
<i>Bauhinia brachycarpa</i> Wall. ex Benth.	-	72
<i>Bauhinia championii</i> (Benth.) Benth.	+	72
<i>Bauhinia hunanensis</i> Hand.Mazz.	+	72
<i>Bauhinia microstachya</i> (Raddi) J.F. Macbr.	-	50
<i>Bauhinia ornata</i> Kurz var. <i>austrosinensis</i> (Tang & Wang) T.C. Chen	-	72
<i>Bauhinia ornata</i> Kurz var. <i>kerri</i> (Gapnep.) K. Larsen & S.S. Larsen	-	72
<i>Bauhinia rufescens</i> Lam.	-	165
<i>Bauhinia scandens</i> L. var. <i>anguina</i> (Roxb.) H. Ohashi	-	76
<i>Bauhinia vahlii</i> Wight & Arn.	+	76
<i>Bauhinia variegata</i> L. var. <i>candida</i> Voigt	-	14
<i>Berlinia bracteosa</i> Benth.	-	77
<i>Berlinia craibiana</i> Baker f.	-	79
<i>Bocoa prouacensis</i> Aubl.	+	15
<i>Bossiaea bossiaeoidea</i> (A. Cunn. ex Benth.) Court	+	135

Taxon	Status ¹	Source ²
<i>Bossiaea foliosa</i> A. Cunn.	+	29
<i>Brachystegia bussei</i> Harms	-	79
<i>Brachystegia cynometroides</i> Harms	-	150
<i>Brachystegia floribunda</i> Benth.	-	79
<i>Brachystegia glaberrima</i> R.E. Fr.	-	79
<i>Brachystegia longifolia</i> Benth.	-	79
<i>Brachystegia spiciformis</i> Benth.	+	69
<i>Brachystegia taxifolia</i> Harms	-	79
<i>Brachystegia wangermeeana</i> De Wild.	-	79
<i>Brownea longipedicellata</i> Huber	-	50
<i>Brya microphylla</i> Bisse	+	150
<i>Bryaspis lupulina</i> (Benth.) P.A. Duvign.	+	42
<i>Caesalpinia cacalaco</i> Bonpl.	-	69
<i>Caesalpinia digyna</i> Rottler	-	76
<i>Caesalpinia eriostachya</i> Benth.	-	44
<i>Caesalpinia pyramidalis</i> Tul.	-	50
<i>Caesalpinia sappan</i> L.	+	72
<i>Caesalpinia velutina</i> (Britton & Rose) Standl.	-	44
<i>Callerya atropurpurea</i> (Wall.) Schot	-	108
<i>Callerya cinerea</i> (Benth.) Schot	+	71
<i>Callerya reticulata</i> (Benth.) Schot	+	71
<i>Callerya speciosa</i> (Champ.) Schot	+	72
<i>Calliandra eriophylla</i> Benth.	+	20
<i>Calliandra harrisii</i> (Lindl.) Benth.	+	50
<i>Calliandra houstoniana</i> (Mill.) Standl. var. <i>calothyra</i> (Meisn.) Barneby	+	146
<i>Calliandra tergemina</i> (L.) Benth. <i>emarginata</i> (Willd.) Barneby	+	5
<i>Calpocalyx dinklagei</i> Harms	-	77
<i>Campylotropis eriocarpa</i> (DC.) Schindl.	+	131
<i>Campylotropis stenocarpa</i> (Klotzsch) Schindl.	+	117
<i>Canavalia brasiliensis</i> Mart. ex Benth.	+	30
<i>Caragana brevispina</i> Royle ex Benth.	+	117
<i>Caragana intermedia</i> Kuang & H.C. Fu	+	155
<i>Caragana leveillei</i> Kom.	-	71
<i>Caragana pleiophylla</i> (Regel) Pojark.	+	72
<i>Caragana polourensis</i> Franch.	+	72
<i>Caragana pruinosa</i> Kom.	+	156
<i>Caragana rosea</i> Turcz. ex Kom.	+	71
<i>Caragana turkestanica</i> Kom.	-	71
<i>Caragana versicolor</i> Benth.	+	169
<i>Cassia montana</i> B. Hayne ex Roth	-	143
<i>Cenostigma tocaninum</i> Ducke	-	50
<i>Centrosema macrocarpum</i> Benth.	-	2
<i>Centrosema sagittatum</i> (Humb. & Bonpl. ex Willd.) Brandegee	+	38
<i>Centrosema schottii</i> (Millsp.) K. Schum.	+	38
<i>Ceratonia siliqua</i> L.	+	82
<i>Cercis chingii</i> Chun	-	71
<i>Cercis gigantea</i> ined.	-	72
<i>Cercis glabra</i> Pamp.	-	71
<i>Chadsia grevei</i> Drake	+	149
<i>Chamaecrista capensis</i> (Thunb.) E. Mey. var. <i>flavescens</i> (E. Mey.) Vogel	+	61
<i>Chamaecrista conferta</i> (Benth.) H.S. Irwin & Barneby	+	152
<i>Chamaecrista conferta</i> (Benth.) H.S. Irwin & Barneby var. <i>virgata</i> (H.S. Irwin & Barneby) H.S. Irwin & Barneby	+	150
<i>Chamaecrista coradinii</i> Barneby	+	152
<i>Chamaecrista decumbens</i> (Benth.) H.S. Irwin & Barneby	+	150
<i>Chamaecrista desvauxii</i> (Collad.) Killip var. <i>brevipes</i> (Benth.) H.S. Irwin & Barneby	+	150
<i>Chamaecrista desvauxii</i> (Collad.) Killip var. <i>circumdatta</i> H.S. Irwin & Barneby	+	150
<i>Chamaecrista desvauxii</i> (Collad.) Killip var. <i>desvauxii</i>	+	84

Taxon	Status ¹	Source ²
<i>Chamaecrista desvauxii</i> (Collad.) Killip var. <i>glauca</i> (Hassl.) H.S. Irwin & Barneby	+	52
<i>Chamaecrista desvauxii</i> (Collad.) Killip var. <i>linearis</i> (H.S. Irwin) H.S. Irwin & Barneby	+	150
<i>Chamaecrista desvauxii</i> (Collad.) Killip var. <i>mollissima</i> (Benth.) H.S. Irwin & Barneby	+	150
<i>Chamaecrista fagonioides</i> (Vogel) H.S. Irwin & Barneby	+	152
<i>Chamaecrista fagonioides</i> (Vogel) H.S. Irwin & Barneby var. <i>macrocalyx</i> (H.S. Irwin & Barneby) H.S. Irwin & Barneby	+	150
<i>Chamaecrista geminata</i> (Benth.) H.S. Irwin & Barneby	+	152
<i>Chamaecrista glandulosa</i> (L.) Greene var. <i>brasiliensis</i> (Vogel) H.S. Irwin & Barneby	+	52
<i>Chamaecrista isidorea</i> (Benth.) H.S. Irwin & Barneby	+	150
<i>Chamaecrista lineata</i> (Sw.) Greene	+	150
<i>Chamaecrista neesiana</i> (Benth.) H. Irwin & Barneby var. <i>subnitida</i> (Taub.) H. Irwin & Barneby	+	150
<i>Chamaecrista nictitans</i> (L.) Moench ssp. <i>patellaria</i> (Coll.) H.S. Irwin & Barneby var. <i>glabrata</i> (Vogel) H.S. Irwin & Barneby	+	68
<i>Chamaecrista orbiculata</i> (Benth.) H.S. Irwin & Barneby	+	152
<i>Chamaecrista orbiculata</i> (Benth.) H.S. Irwin & Barneby var. <i>orbiculata</i>	+	150
<i>Chamaecrista pilosa</i> (L.) Greene var. <i>pilosa</i>	+	24
<i>Chamaecrista plumosa</i> (E. Mey) H. Irwin & Barneby. var. <i>plumosa</i>	+	61
<i>Chamaecrista ramosa</i> (Vogel) H.S. Irwin & Barneby var. <i>curvifolia</i> (Vogel) H.S. Irwin & Barneby	+	150
<i>Chamaecrista rotundifolia</i> (Pers.) Greene var. <i>rotundifolia</i>	+	150
<i>Chamaecrista serpens</i> (L.) Greene var. <i>serpens</i>	+	150
<i>Chamaecrista serpens</i> (L.) Greene var. <i>wrightii</i> (A. Gray) H.S. Irwin & Barneby	+	68
<i>Chamaecrista tetraphila</i> Desv.	+	84
<i>Chesneya cuneata</i> (Benth.) Ali	-	117
<i>Cladrastis wilsonii</i> Takeda	+	71
<i>Clitoria sagotii</i> Fantz	+	150
<i>Cojoba arborea</i> (L.) Britton & Rose var. <i>arborea</i>	+	68
<i>Colutea istria</i> Mill.	+	44
<i>Conzattia multiflora</i> (B.L. Rob.) Standl.	-	150
<i>Craibia brevicaudata</i> (Vatke) Dunn	+	69
<i>Cratylia hypargyrea</i> Mart.	+	50
<i>Crotalaria angulata</i> Mill.	+	76
<i>Crotalaria atrorubens</i> Hochst. ex Benth.	+	49
<i>Crotalaria brevidens</i> Benth.	+	38
<i>Crotalaria brevis</i> Domin	+	65
<i>Crotalaria chinensis</i> L.	+	57
<i>Crotalaria eremaea</i> F. Muell. ssp. <i>eremaea</i>	+	65
<i>Crotalaria excisa</i> (Thunb.) Baker f.	+	60
<i>Crotalaria ferruginea</i> Graham ex Benth.	+	76
<i>Crotalaria glaucoides</i> Baker f.	+	42
<i>Crotalaria hainanensis</i> Huang	+	57
<i>Crotalaria humifusa</i> Benth.	+	76
<i>Crotalaria humilis</i> Eckl. & Zeyh.	+	60
<i>Crotalaria lathyroides</i> Guill. & Perr.	+	42
<i>Crotalaria mitchellii</i> Benth.	+	65
<i>Crotalaria perrottetii</i> DC.	+	140
<i>Crotalaria saharae</i> Coss.	+	150
<i>Crotalaria sessiliflora</i> L. ssp. <i>hazarensis</i> Ali	+	9
<i>Crotalaria sessiliflora</i> L. ssp. <i>sessiliflora</i>	+	9
<i>Crotalaria trifoliatum</i> Willd.	+	146
<i>Cryptosepalum exfoliatum</i> De Wild.	-	79
<i>Cullen australasicum</i> (Schltdl.) J.W. Grimes	+	65
<i>Cullen cinereum</i> (Lindl.) J.W. Grimes	+	65
<i>Cyathostegia mathewsii</i> (Benth.) Schery	+	150
<i>Cyclolobium vecchii</i> A. Samp. ex Hoehne	+	50

Taxon	Status ¹	Source ²
<i>Cyclopia buxifolia</i> (Burm. f.) Kies	+	153
<i>Cyclopia dregeana</i> Kies	+	153
<i>Cyclopia intermedia</i> E. Mey.	+	153
<i>Cyclopia latifolia</i> DC.	+	153
<i>Cyclopia longifolia</i> Vogel	+	153
<i>Cyclopia meyeriana</i> Walp.	+	153
<i>Cyclopia pubescens</i> Eckl. & Zeyh.	+	153
<i>Cyclopia sessiliflora</i> Eckl. & Zeyh.	+	153
<i>Cyclopia subternata</i> Vogel	+	153
<i>Cylicodiscus gabunensis</i> Harms	-	150
<i>Cytisus arboreus</i> (Desf.) DC.	+	1
<i>Dahlstedtia pinnata</i> (Benth.) Malme	-	50
<i>Dalbergia albiflora</i> A. Chev. ex Hutch. & Dalziel	+	99
<i>Dalbergia balansae</i> Prain	+	72
<i>Dalbergia brasiliense</i> Vogel	+	50
<i>Dalbergia candenatensis</i> (Dennst.) Prain	+	38
<i>Dalbergia cultrata</i> Benth.	+	72
<i>Dalbergia dalzielii</i> Baker f. ex Hutch. & Dalziel	+	99
<i>Dalbergia decipularis</i> Rizzini & A. Mattos	+	50
<i>Dalbergia dyeriana</i> Prain	-	71
<i>Dalbergia hainanensis</i> Merr. & Chun	+	72
<i>Dalbergia hancei</i> Benth.	+	71
<i>Dalbergia hostilis</i> Benth.	+	99
<i>Dalbergia hupeana</i> Hance	+	72
<i>Dalbergia lanceolaria</i> L. f. var. <i>assamica</i> (Benth.) Thoth.	+	72
<i>Dalbergia miscolobium</i> Benth.	+	50
<i>Dalbergia obtusifolia</i> (Baker) Prain	+	72
<i>Dalbergia odorifera</i> T.C. Chen	+	72
<i>Dalbergia oligophylla</i> Hutch. & Dalziel	+	150
<i>Dalbergia pinnata</i> (Lour.) Prain	+	72
<i>Dalbergia riparia</i> (Mart.) Benth.	+	50
<i>Dalbergia rufa</i> G. Don	+	99
<i>Dalbergia saxatilis</i> Hook. f.	+	99
<i>Dalbergia sericea</i> G. Don	+	76
<i>Dalbergia stipulacea</i> Roxb.	+	76
<i>Dalbergia tsoi</i> Merr. & Chun	+	72
<i>Dalbergia villosa</i> (Benth.) Benth.	+	50
<i>Dalea mollissima</i> (Rydb.) Munz	+	48
<i>Dalea virgata</i> Lag.	+	150
<i>Daniellia alsteeniana</i> Duvign.	-	79
<i>Daniellia oliveri</i> (Rolfe) Hutch. & Dalziel	-	165
<i>Daviesia decurrens</i> Meisn.	+	151
<i>Daviesia horrida</i> L. Preiss ex Meisn.	+	149
<i>Daviesia mimosoides</i> R. Br.	+	114
<i>Dendrolobium dispernum</i> (Hayata) Schindl.	+	150
<i>Dendrolobium triangulare</i> (Retz.) Schindl.	+	72
<i>Derris cumingii</i> Benth.	+	43
<i>Derris cuneifolia</i> Benth.	+	76
<i>Derris marginata</i> (Roxb.) Benth.	+	76
<i>Derris trifoliata</i> Lour.	+	72
<i>Desmanthus velutinus</i> Scheele	+	146
<i>Desmodium axillare</i> (Sw.) DC.	+	38
<i>Desmodium campylocaulon</i> F. Muell. ex Benth.	+	38
<i>Desmodium concinnum</i> DC.	+	76
<i>Desmodium confertum</i> DC.	+	76
<i>Desmodium cuneatum</i> Hook. & Arn.	+	38
<i>Desmodium flagellare</i> Benth.	+	65
<i>Desmodium laburnifolium</i> (Poir.) DC.	+	143

Taxon	Status ¹	Source ²
<i>Desmodium macrocarpum</i> Domin	+	65
<i>Desmodium multiflorum</i> DC.	+	131
<i>Desmodium neomexicanus</i> A. Gray	+	146
<i>Desmodium podocarpum</i> DC.	+	9
<i>Desmodium podocarpum</i> DC. var. <i>szechuenense</i> Craib	+	72
<i>Desmodium renifolium</i> (L.) Schindl.	+	76
<i>Desmodium rubrum</i> (Lour.) DC.	+	72
<i>Desmodium salicifolium</i> (Poir.) DC.	+	24
<i>Desmodium sandwicense</i> E. Mey.	+	146
<i>Desmodium sinuatum</i> Baker	+	155
<i>Desmodium trichostachyum</i> Benth.	+	65
<i>Detarium macrocarpum</i> Harms	-	150
<i>Detarium senegalense</i> J.F. Gmel.	-	165
<i>Dicorynia guianensis</i> Amshoff	+	15
<i>Dicorynia guianensis</i> Amshoff	-	62
<i>Didelotia africana</i> Baill.	-	77
<i>Dillwynia floribunda</i> Sm. var. <i>floribunda</i>	+	65
<i>Dillwynia sericea</i> A. Cunn.	+	106
<i>Dimorphandra conjugata</i> (Splitg.) Sandwith	+	150
<i>Dimorphandra vernicosa</i> Benth.	+	50
<i>Diplostropis purpurea</i> (Rich.) Amshoff var. <i>purpurea</i>	+	150
<i>Discolobium leptophyllum</i> Benth.	+	88
<i>Discolobium psoraliifolium</i> Benth.	+	109
<i>Discolobium pulchellum</i> Benth.	+	109
<i>Dorycnium pentaphyllum</i> Scop. ssp. <i>germanicum</i> (Gremli) Gams	+	112
<i>Duparquetia orchidacea</i> Baill.	-	150
<i>Dussia martinicensis</i> Krug & Urb. ex Taub.	+	142
<i>Dussia martinicensis</i> Krug & Urb. ex Taub.	-	150
<i>Ebenopsis confinis</i> (Standl.) Barneby & J.W. Grimes	+	150
<i>Ebenus pinnata</i> Aiton	+	150
<i>Elizabetha durissima</i> Ducke	-	50
<i>Elizabetha paraensis</i> Ducke	-	50
<i>Elizabetha princeps</i> Schomb. ex Benth.	-	50
<i>Elizabetha speciosa</i> Ducke	-	50
<i>Entada africana</i> Guill. & Perr.	+	69
<i>Entada mannii</i> (Oliv.) Tisser.	+	150
<i>Entada rheedei</i> Spreng. ssp. <i>sinohimalensis</i> (Grierson & Long) Panigr.	+	76
<i>Eriosema ellipticifolium</i> Schinz	+	61
<i>Eriosema glomeratum</i> (Guill. & Perr.) Hook. f.	+	42
<i>Erythrina americana</i> Mill.	+	69
<i>Erythrina arborescens</i> Roxb.	+	76
<i>Erythrina barqueroana</i> Krukoff & Barneby	+	50
<i>Erythrina brucei</i> Schweinf.	+	8
<i>Erythrina cochleata</i> Stnagl.	+	44
<i>Erythrina edulis</i> Triana	+	137
<i>Erythrina sandwicensis</i> O. Deg.	+	44
<i>Erythrina senegalensis</i> DC.	+	165
<i>Erythrophleum chlorostachys</i> (F. Muell.) Baill.	+	150
<i>Erythrophleum fordii</i> Oliv.	+	41
<i>Erythrophleum ivorense</i> A. Chev.	+	77
<i>Exostyles venusta</i> Spreng.	-	50
<i>Eysenhardtia polystachya</i> (Ortega) Sarg.	-	68
<i>Eysenhardtia texana</i> Scheele	+	146
<i>Flemingia procumbens</i> Roxb.	+	131
<i>Flemingia prostrata</i> Roxb.	+	150
<i>Galactia rubra</i> (Jacq.) Urb.	+	150
<i>Gastrolobium bilobium</i> R. Br.	+	151
<i>Genista falcata</i> Brot.	+	60

Taxon	Status ¹	Source ²
<i>Genista ferox</i> Poir.	+	150
<i>Genista microcephala</i> Coss. & Durieru	+	150
<i>Genista numidica</i> Spach ssp. <i>numidica</i>	+	150
<i>Genista tinctoria</i> L. var. <i>ovata</i> (Waldst. & Kit) F.W. Schultz	+	5
<i>Genista tricuspidata</i> Desf.	+	150
<i>Gilbertiodendron mayombense</i> (Pellegr.) J. L. onard	-	77
<i>Gleditsia amorphoides</i> (Griseb.) Taub.	-	6
<i>Gleditsia anstralis</i> Hemsl.	-	71
<i>Gleditsia japonica</i> Miq. var. <i>japonica</i>	-	73
<i>Gleditsia japonica</i> Miq. var. <i>velutina</i> L.C. Li	-	71
<i>Gleditsia macrantha</i> Desf.	+	130
<i>Gliricidia sepium</i> (Jacq.) Kunth ex Walp.	-	108
<i>Glycine falcata</i> Benth.	+	65
<i>Glycine microphylla</i> (Benth.) Tindale	+	75
<i>Glycine soja</i> Siebold & Zucc.	+	146
<i>Glycyrrhiza pallidiflora</i> Maxim.	+	32
<i>Gneldenstaedtia himalaica</i> Baker	+	169
<i>Gneldenstaedtia multiflora</i> Bunge	+	155
<i>Gneldenstaedtia verna</i> (Georgi) Boriss.	+	156
<i>Gneldenstaedtia verna</i> (Georgi) Boriss. ssp. <i>multiflora</i> (Bunge) H.P. Tsui	+	156
<i>Gymnocladus chinensis</i> Baill.	-	72
<i>Harpalyce cubensis</i> Griseb.	+	150
<i>Harpalyce macrocarpa</i> Britton & P. Wilson	+	150
<i>Hedysarum aculeolatum</i> Boiss.	+	150
<i>Hedysarum carnosum</i> Desf.	+	146
<i>Hedysarum flexosum</i> L.	+	146
<i>Hedysarum humile</i> L.	+	150
<i>Hedysarum pallens</i> Desf.	+	95
<i>Hedysarum pallidum</i> Desf.	+	150
<i>Hesperalbizia occidentalis</i> (Brandege) Barbeny & J.W. Grimes	+	150
<i>Hippocrepis minor</i> Munby	+	150
<i>Humboldtia laurifolia</i> Vahl	+	164
<i>Hymenolobium janeirensense</i> Kuhlmann	-	50
<i>Hymenostegia afzelii</i> (Oliv.) Harms	-	77
<i>Hypocalyptus coluteoides</i> (Lam.) R. Dahlgren	+	150
<i>Indigastrum parviflorum</i> (Heyne ex Wight & Arn.) Schrire	+	65
<i>Indigofera amoena</i> Aiton	+	60
<i>Indigofera aquae-nitensis</i> Bremek.	+	60
<i>Indigofera argyraea</i> Eckl. & Zeyh.	+	60
<i>Indigofera brevidens</i> Benth.	+	167
<i>Indigofera bungeana</i> Walp.	+	71
<i>Indigofera candicans</i> Aiton	+	60
<i>Indigofera capillaris</i> Thunb.	+	60
<i>Indigofera chuniana</i> F.P. Metcalf	+	72
<i>Indigofera coriacea</i> Aiton var. <i>coriacea</i>	+	61
<i>Indigofera coriacea</i> Aiton var. <i>minor</i> (E. Mey.) Harv.	+	61
<i>Indigofera cylindracea</i> Wall. ex Baker	+	76
<i>Indigofera daleoides</i> Benth. ex Harv. var. <i>daleoides</i>	+	61
<i>Indigofera filifolia</i> Thunb.	+	61
<i>Indigofera flabellata</i> Harv.	+	61
<i>Indigofera fortunei</i> Craib	+	71
<i>Indigofera galeoides</i> DC.	+	72
<i>Indigofera hebeptala</i> Benth. Ex Baker	+	76
<i>Indigofera heterantha</i> Brandis	+	117
<i>Indigofera incana</i> Thunb.	+	61
<i>Indigofera lespedezioides</i> Kunth	-	83
<i>Indigofera linmaei</i> Ali	+	38
<i>Indigofera litoralis</i> Chun & T.C. Chen	+	57

Taxon	Status ¹	Source ²
<i>Indigofera miniata</i> Ortega var. <i>leptosepala</i> (Nutt. ex Torr. A. Gray) B.L. Turner	+	146
<i>Indigofera neglectum</i> Ledeb.	+	146
<i>Indigofera paniculata</i> Pers.	+	49
<i>Indigofera pratensis</i> F. Muell.	+	65
<i>Indigofera pulchra</i> Willd.	+	49
<i>Indigofera rhodantha</i> Fourc.	+	60
<i>Indigofera senegalensis</i> Lam.	+	42
<i>Indigofera simplicifolia</i> Lam.	+	49
<i>Indigofera trifoliata</i> L.	+	38
<i>Indigofera hendecaphylla</i> Jacq.	+	146
<i>Indopiptadenia oudhensis</i> (Brandis) Brenan	+	76
<i>Inga chocoensis</i> Killip ex T.S. Elias	+	150
<i>Inga congesta</i> T.D. Penn.	+	50
<i>Inga cordistipula</i> Mart.	+	50
<i>Inga hispida</i> Schott ex Benth.	+	50
<i>Inga jinicuil</i> G. Don	+	69
<i>Inga lenticifolia</i> Benth.	+	50
<i>Inga maritima</i> Benth.	+	50
<i>Inga nobilis</i> Willd. ssp. <i>quaternata</i> (Poepp. & Endl.) T.D. Penn.	+	68
<i>Inga oerstedia</i> Benth.	+	69
<i>Inga semialata</i> (Vell.) Mart.	+	136
<i>Isoberlinia angolensis</i> (Benth.) Hoyle & Brenan	-	79
<i>Isotropis atropurpurea</i> F. Muell.	+	167
<i>Jacksonia dilatata</i> Benth.	+	135
<i>Julbernardia paniculata</i> (Benth.) Troupin	-	79
<i>Koompassia malaccensis</i> Maingay ex Benth.	-	69
<i>Lablab pourpureus</i> (L.) Sweet subsp. <i>Purpureus</i>	+	5
<i>Lathyrus humilis</i> (Ser.) Fischer ex Spreng.	+	9
<i>Lathyrus lanszwertii</i> Kellogg var. <i>leucanthus</i> (Rydb.) Dorn	+	91
<i>Lathyrus littoralis</i> (Nutt.) Endl.	+	80
<i>Lathyrus szowitsii</i> Boiss.	+	27
<i>Lebeckia multiflora</i> E. Mey.	+	60
<i>Leptoderris nobilis</i> (Bajer) Dunn	+	150
<i>Lespedeza buergeri</i> Miq.	+	72
<i>Lespedeza chinensis</i> G. Don	+	72
<i>Lespedeza daurica</i> (Laxm.) Schindl	+	146
<i>Lespedeza</i> × <i>divaricata</i> (Nakai) T.B. Lee	+	150
<i>Lespedeza floribunda</i> Bunge	+	72
<i>Lespedeza formosa</i> (Vogel) Koehne ssp. <i>elliptica</i> (Benth. ex Maxim.) S. Akiyama & H. Ohba	+	72
<i>Lespedeza homoloba</i> Nakai	+	61
<i>Lespedeza japonica</i> L.H. Bailey	+	146
<i>Lespedeza tomentosa</i> (Thunb.) Siebold ex Maxim.	+	146
<i>Lessertia brachypus</i> Harv.	+	60
<i>Lessertia capitata</i> E. Mey.	+	60
<i>Lessertia diffusa</i> R. Br.	+	60
<i>Lessertia excisa</i> DC.	+	60
<i>Lessertia inflata</i> Harv.	+	61
<i>Lessertia macrostachya</i> DC.	+	60
<i>Lessertia spinescens</i> E. Mey.	+	60
<i>Leucaena collinsii</i> Britton & Rose	+	69
<i>Leucaena collinsii</i> Britton & Rose ssp. <i>collinsii</i>	+	116
<i>Leucaena collinsii</i> Britton & Rose ssp. <i>zacapana</i> C.E. Hughes	+	116
<i>Leucaena cuspidata</i> Standl.	+	116
<i>Leucaena esculenta</i> (DC.) Benth. ssp. <i>esculenta</i>	+	153
<i>Leucaena esculenta</i> (DC.) Benth. ssp. <i>matudae</i> Z rate	+	153
<i>Leucaena greggii</i> S. Watson	+	116
<i>Leucaena involucrata</i> Z rate	+	116

Taxon	Status ¹	Source ²
<i>Leucaena lanceolata</i> S. Watson	+	69
<i>Leucaena lanceolata</i> S. Watson ssp. <i>lanceolata</i>	+	116
<i>Leucaena lanceolata</i> S. Watson ssp. <i>sousae</i> Z rate	+	116
<i>Leucaena lanceolata</i> S. Watson var. <i>lanceolata</i>	+	116
<i>Leucaena lanceolata</i> S. Watson var. <i>sousae</i> (Z rate) C.E. Hughes	+	116
<i>Leucaena lempirana</i> C.E. Hughes	+	116
<i>Leucaena macrophylla</i> Benth.	+	69
<i>Leucaena macrophylla</i> Benth. ssp. <i>macrophylla</i>	+	116
<i>Leucaena macrophylla</i> Benth. ssp. <i>nelsonii</i> (Britton & Rose) Z rate	+	116
<i>Leucaena magnifica</i> (C.E. Hughes) C.E. Hughes	+	116
<i>Leucaena matudae</i> (Z rate) C.E. Hughes	+	116
<i>Leucaena multicapitula</i> Schery	+	116
<i>Leucaena pallida</i> Britton & Rose	+	20
<i>Leucaena pueblana</i> Britton & Rose	+	116
<i>Leucaena retusa</i> Benth.	+	69
<i>Leucaena retusa</i> Benth.	-	70
<i>Leucaena salvadorensis</i> Standl. ex Britton & Rose	+	76
<i>Leucaena shannonii</i> Donn. Sm.	+	69
<i>Leucaena shannonii</i> Donn. Sm. ssp. <i>magnifica</i> C.E. Hughes	+	116
<i>Leucaena shannonii</i> Donn. Sm. ssp. <i>shannonii</i>	+	116
<i>Leucaena trichandra</i> (Zucc.) Urb.	+	116
<i>Leucomphalos capparideus</i> Planch.	-	150
<i>Liparia parva</i> Vogel ex Walp.	+	61
<i>Liparia splendens</i> (Burm. f.) Bos & De Wit ssp. <i>splendens</i>	+	61
<i>Loesenera talbotii</i> Baker f.	-	150
<i>Lonchocarpus filipes</i> Benth.	+	50
<i>Lonchocarpus leucanthus</i> Burkart	+	50
<i>Lonchocarpus muehlbergianus</i> Hassl.	+	50
<i>Lonchocarpus roseus</i> (Mill.) DC.	+	142
<i>Lonchocarpus spruceanus</i> Benth.	+	50
<i>Lotononis benthamiana</i> Dumm.	+	60
<i>Lotononis brachyloba</i> Benth.	+	60
<i>Lotononis digitata</i> Harv.	+	61
<i>Lotononis falcata</i> (E. Mey.) Benth.	+	60
<i>Lotononis foliosa</i> H. Bol.	+	60
<i>Lotononis lenticula</i> (E. Mey.) Benth.	+	61
<i>Lotononis leptoloba</i> H. Bol.	+	60
<i>Lotononis longiflora</i> H. Bol.	+	60
<i>Lotononis magnistipulata</i> Dumm.	+	60
<i>Lotononis maximiliani</i> De Wild.	+	60
<i>Lotononis pentaphylla</i> Benth.	+	60
<i>Lotononis quinata</i> Benth.	+	60
<i>Lotononis serpens</i> (E. Mey.) Dahlgren	+	61
<i>Lotononis strigillosa</i> (Merxm. & A. Schreib.) A. Schreib.	+	61
<i>Lotus alpinus</i> (DC.) Schleich. ex Ramond	+	85
<i>Lotus corniculatus</i> L. var. <i>hirsutus</i> W.D.J. Kock	+	112
<i>Lotus kyylovii</i> Schischk. & Serg.	+	125
<i>Lotus macrotrichus</i> Boiss.	+	89
<i>Lotus mearnsii</i> (Britton) Greene	+	58
<i>Lotus preslii</i> Ten.	+	66
<i>Lotus rigidus</i> (Benth.) Greene	+	48
<i>Lotus salsuginosus</i> Greene	+	146
<i>Lotus schoelleri</i> Schweinf.	+	125
<i>Lotus strigosus</i> (Nutt.) Greene var. <i>tomentellus</i> (Greene) Isely	+	48
<i>Luetzelburgia auriculata</i> (Allemao) Ducke	-	50
<i>Luetzelburgia guaissara</i> Toledo	-	50
<i>Lupinus argenteus</i> Pursh	+	146
<i>Lupinus argenteus</i> Pursh var. <i>holosericeus</i> (Nutt.) Barneby	+	92

Taxon	Status ¹	Source ²
<i>Lupinus argenteus</i> Pursh var. <i>rubricaulis</i> (Greene) S.L. Welsh	+	91
<i>Lupinus atlanticus</i> Gladst.	+	146
<i>Lupinus campestris</i> Schltdl. & Cham.	+	12
<i>Lupinus chamissonis</i> Eschsch.	+	80
<i>Lupinus cosentinii</i> Guss.	+	128
<i>Lupinus exaltatus</i> Zucc.	+	12
<i>Lupinus formosus</i> Greene	+	146
<i>Lupinus hispanicus</i> Boiss. & Reut.	+	146
<i>Lupinus meridanus</i> Moritz ex C.P. Sm.	+	160
<i>Lupinus montanus</i> Kunth	+	12
<i>Lupinus princei</i> Harms	+	81
<i>Lupinus sericeus</i> Pursh	+	146
<i>Lupinus texensis</i> Hook.	+	5
<i>Lysiloma divaricatum</i> (Jacq.) J.F. Macbr.	+	150
<i>Lysiloma latisiliquum</i> (L.) Benth.	+	118
<i>Lysiloma tergeminum</i> Benth.	+	150
<i>Machaerium angustifolium</i> Vogel	+	50
<i>Machaerium arboreum</i> (Jacq.) Vogel	+	126
<i>Machaerium brasiliense</i> Vogel	+	50
<i>Machaerium gracile</i> Benth.	+	50
<i>Machaerium hirtum</i> (Vell.) Stellfield	+	50
<i>Machaerium milleflorum</i> Pittier	+	126
<i>Machaerium myrianthum</i> Benth.	+	150
<i>Machaerium quinatum</i> (Aubl.) Sandwith var. <i>quinatum</i>	+	150
<i>Machaerium stipitatum</i> (DC.) Vogel	+	50
<i>Machaerium triste</i> Vogel	+	50
<i>Macroptilium gibbosifolium</i> (Ortega) A. Delgado	+	146
<i>Macroptilium gracile</i> (Poepp. ex Benth.) Urb.	+	38
<i>Macrotyloma uniflorum</i> (Lam.) Verdc.	+	146
<i>Medicago archiducis-nicolai</i> Sirj.	+	169
<i>Medicago doliata</i> Carmign.	+	11
<i>Medicago granadensis</i> Willd.	+	22
<i>Medicago noeana</i> Boiss.	+	119
<i>Medicago polyceratia</i> (L.) Trautv.	+	86
<i>Medicago radiata</i> L.	+	42
<i>Medicago rigiduloides</i> E. Small	+	11
<i>Medicago rotata</i> Boiss.	+	146
<i>Medicago sativa</i> L. nothosp. <i>Varia</i> (Martyn) Arcang.	+	147
<i>Meizotropis buteiformis</i> Voigt	+	76
<i>Melilotus infestus</i> Guss.	+	150
<i>Melilotus macrocarpus</i> Coss. & Durieru	+	150
<i>Melilotus officinalis</i> Lam.	+	5
<i>Melolobium aethiopicum</i> (L.) Druce	+	61
<i>Melolobium calycinum</i> Benth.	+	60
<i>Melolobium humile</i> Eckl. & Zeyh.	+	60
<i>Microberlinia bisulcata</i> A. Chev.	-	150
<i>Mildbraediodendron excelsum</i> Harms	-	150
<i>Millettia extensa</i> (Benth.) Benth. ex Baker	+	76
<i>Millettia ferruginea</i> (Hochst.) Baker	+	67
<i>Millettia glaucescens</i> Kurz	+	76
<i>Millettia laurentii</i> De Wild.	+	93
<i>Millettia thonningii</i> (Schumach.) Baker	+	69
<i>Millettia usaramoensis</i> Taub.	+	5
<i>Mimosa affinis</i> B.L. Rob.	+	161
<i>Mimosa arenosa</i> (Willd.) Poir. Var. <i>arenosa</i>	+	68
<i>Mimosa artemisiana</i> Heringer & Paula	+	50
<i>Mimosa ceratonia</i> L.	+	68
<i>Mimosa laticifera</i> Rizzini & A. Mattos	+	50

Taxon	Status ¹	Source ²
<i>Mimosa luisana</i> Brandegees	+	150
<i>Mimosa ophthalmocentra</i> Mart. ex Benth.	+	150
<i>Mimosa polyantha</i> Benth.	+	150
<i>Mimosa quadrivalvis</i> L. var. <i>nutallii</i> (DC.) Beard	+	146
<i>Mimosa rubicaulis</i> Lam. ssp. <i>himalayana</i> (Gamble) H. Ohashi	+	76
<i>Mucuna bracteata</i> DC. ex Kurz	+	97
<i>Mucuna sempervirens</i> Hemsl.	+	71
<i>Neonotonia wightii</i> (Wight & Arn) J.A. Lackey	+	38
<i>Neptunia pubescens</i> Benth.	+	88
<i>Newtonia hildebrandtii</i> (Vatke) Torre	-	68
<i>Omosia krugii</i> Urb.	+	68
<i>Onobrychis kabylica</i> (Bornm.) Sirj.	+	150
<i>Onobrychis paucidentata</i> Pomel	+	150
<i>Ononis alba</i> Poir.	+	150
<i>Ononis antennata</i> Pomel ssp. <i>antennata</i>	+	150
<i>Ononis euphrasiifolia</i> Desf.	+	150
<i>Ormocarpum sennoides</i> (Willd.) DC.	+	50
<i>Ormosia apiculata</i> L. Chen	+	72
<i>Ormosia emarginata</i> (Hook. & Arn.) Benth.	+	72
<i>Ormosia glauca</i> Wall.	+	76
<i>Ormosia henryi</i> Prain	+	72
<i>Ormosia hosei</i> Hemsl. & E.H. Wilson	+	71
<i>Ormosia pinnata</i> (Lour.) Merr.	+	41
<i>Ormosia semicastrata</i> Hance	+	72
<i>Ormosia semicastrata</i> Hance f. <i>litchiifolia</i> F.C. How	+	72
<i>Ormosia semicastrata</i> Hance f. <i>pallida</i> F.C. How	+	72
<i>Ormosia xylocarpa</i> Chun ex Merr. & L. Chen	+	71
<i>Otholobium glandulosum</i> (L.) J.W. Grimes	+	38
<i>Otholobium striatum</i> (Thunb.) C.H. Stirt.	+	60
<i>Oxylobium scandens</i> (Sm.) Benth.	+	150
<i>Oxyrhynchus volubilis</i> Brandegees	+	149
<i>Oxytropis anertii</i> Nakai	+	150
<i>Oxytropis campanulata</i> Vassilcz.	+	113
<i>Oxytropis kansuensis</i> Bunge	+	169
<i>Oxytropis melanocalyx</i> Bunge	+	169
<i>Oxytropis nigrescens</i> (Pall.) Fisch. ex DC. var. <i>uniflora</i> (Hook.) Barneby	+	133
<i>Oxytropis sericea</i> Nutt.	+	27
<i>Oxytropis splendens</i> Douglas ex Hook.	+	101
<i>Pachyrhizus ahipa</i> (Wedd.) Parodi	+	96
<i>Pachyrhizus tuberosus</i> (Lam.) Spreng.	+	146
<i>Paramacrolobium coeruleum</i> (Taub.) J. Léonard	-	69
<i>Parapiptadenia excelsa</i> (Griseb.) Burkart	+	6
<i>Paraserianthes toona</i> (F.M. Bailey) I.C. Nielsen	+	20
<i>Parkia bicolor</i> A. Chev.	-	150
<i>Parkia biglobosa</i> (Jacq.) R. Br. ex G. Don	-	165
<i>Parkia filicoidea</i> Welw. ex Oliv.	+	20
<i>Parkia ulei</i> (Harms) Kuhlman. var. <i>surinamensis</i> Kleinhoonte	-	150
<i>Parkinsonia aculeata</i> L.	+	68
<i>Parkinsonia microphylla</i> Torr.	-	26
<i>Peltophorum dasyrhachis</i> (Miq.) Kurz	-	139
<i>Pentaclethra macrophylla</i> Benth.	+	44
<i>Phaseolus coccineus</i> L. subsp. <i>coccineus</i>	+	5
<i>Phaseolus coccineus</i> L. var. <i>darwinianus</i> Hern.-Xol. & Miranda-Colín	+	146
<i>Phaseolus filiformis</i> Benth.	+	146
<i>Piptadeniastrum africanum</i> (hook. f.) Brenan	+	93
<i>Piscidia grandifolia</i> (Donn. Sm.) I.M. Johnst.	+	150
<i>Pithecellobium diversifolium</i> Benth.	+	50
<i>Plathymenia foliolosa</i> Benth.	+	154

Taxon	Status ¹	Source ²
<i>Podalyria burchellii</i> DC.	+	60
<i>Podalyria canescens</i> E. Mey.	+	61
<i>Podalyria cordata</i> R. Br.	+	61
<i>Podalyria velutina</i> Benth.	+	60
<i>Poitea punicea</i> (Urb.) Lavin	+	68
<i>Pongamia pinnata</i> (L.) Pierre	-	108
<i>Prosopis affinis</i> Spreng.	+	56
<i>Prosopis africana</i> (Guille. et al.) Taub.	+	69
<i>Prosopis alba</i> Griseb.	+	69
<i>Prosopis articulata</i> S. Watson	+	69
<i>Prosopis ferox</i> Griseb.	+	6
<i>Prosopis flexuosa</i> DC.	+	6
<i>Prosopis glandulosa</i> Torr. var. <i>glandulosa</i>	+	54
<i>Prosopis glandulosa</i> Torr. var. <i>torreyana</i> (L.D. Benson) M.C. Johnst.	+	48
<i>Prosopis kuntzei</i> Harms	+	69
<i>Prosopis pallida</i> (Humb. & Bonpl. ex Willd.) Kunth	+	69
<i>Prosopis ruscifolia</i> Griseb.	+	69
<i>Prosopis tamarugo</i> F. Phil.	+	69
<i>Prosopis velutina</i> Wooton	+	54
<i>Psoralea aculeata</i> L.	+	61
<i>Psoralea affinis</i> Eckl. & Zeyh.	+	61
<i>Psoralea alata</i> (Thunb.) T.M. Salter	+	61
<i>Psoralea verrucosa</i> Willd.	+	61
<i>Psoralea emoryi</i> (A. Gray) Rydb.	+	48
<i>Psoralea schottii</i> (Torr.) Barneby	+	48
<i>Pterocarpus macrocarpus</i> Kurz	+	148
<i>Pterocarpus osun</i> Craib	+	150
<i>Pterocarpus santalinus</i> L. f.	+	143
<i>Pueraria montana</i> (Lour.) Merr. var. <i>montana</i>	+	5
<i>Pueraria montana</i> (Lour.) Merr. var. <i>thomsonii</i> (Benth.) Wiersema ex D.B. Ward	+	72
<i>Pueraria wallichii</i> DC.	+	76
<i>Pycnospora lutescens</i> (Poir.) Schindl.	+	65
<i>Racosperma oshanesii</i> (F. Muell. & Maiden) Pedley	+	63
<i>Rafnia axillaris</i> Thunb.	+	61
<i>Rafnia capensis</i> (L.) Druce	+	61
<i>Rafnia racemosa</i> Eckl. & Zeyh.	+	61
<i>Recordoxylon speciosum</i> (Benoist) Gazel ex Barneby	+	62
<i>Rhynchosia cunninghamii</i> Benth.	+	24
<i>Rhynchosia harveyi</i> Eckl. & Zeyh.	+	61
<i>Rhynchosia himalensis</i> Benth. ex Baker	+	9
<i>Rhynchosia hirsuta</i> Eckl. & Zeyh.	+	60
<i>Rhynchosia namaensis</i> Schinz	+	60
<i>Rothia indica</i> (L.) Druce	+	16
<i>Samanea inopinata</i> (Harms) Barneby & J.W. Grimes	+	50
<i>Schizolobium parabyba</i> (Vell.) S.F. Macbr.	+	68
<i>Schleinitzia insularum</i> (Guill.) Burkart	+	38
<i>Sclerolobium guianensis</i> Benth.	+	136
<i>Sclerolobium melinonii</i> Harms	+	62
<i>Sclerolobium pilgerianum</i> Harms	+	50
<i>Senna australis</i> (Vell.) H.S. Irwin & Barneby	-	50
<i>Senna aversiflora</i> (Herb.) H.S. Irwin & Barneby	-	50
<i>Senna barclayana</i> (Sweet) Randell var. <i>barclayana</i>	+	88
<i>Senna cumingii</i> (Hook. & Arn.) H.S. Irwin & Barneby	-	6
<i>Senna lindheimeriana</i> (Scheele) H.S. Irwin & Barneby	-	6
<i>Senna obtusifolia</i> (L.) H.S. Irwin & Barneby	+	132
<i>Senna septentrionalis</i> (Viv.) H.S. Irwin & Barneby	+	72
<i>Senna silvestris</i> (Vell.) H.S. Irwin & Barneby ssp. <i>silvestris</i> var. <i>sapindifolia</i> (Vogel) H.S. Irwin & Barneby	-	50

Taxon	Status ¹	Source ²
<i>Senna tora</i> (L.) Roxb.	+	107
<i>Sesbania benthamiana</i> Domin	+	65
<i>Sesbania brachycarpa</i> F. Muell.	+	65
<i>Sesbania emerus</i> (Aubl.) Urb.	+	142
<i>Sesbania erubescens</i> (Benth.) N.T. Burb.	+	65
<i>Sesbania exasperata</i> Kunth	+	51
<i>Sesbania goetzei</i> Harms	+	110
<i>Sesbania leptocarpa</i> DC.	+	49
<i>Sesbania macroptera</i> Micheli	-	6
<i>Sesbania pachycarpa</i> DC.	+	42
<i>Sesbania sesban</i> (L.) Merr. ssp. <i>punctata</i> (DC.) J.B. Gillett	+	141
<i>Sesbania sesban</i> (L.) Merr. ssp. <i>sesban</i>	+	111
<i>Shutteria densiflora</i> Benth.	+	131
<i>Shutteria involucrata</i> (Wall.) Wight & Arn. var. <i>involucrata</i>	-	71
<i>Sindora siamensis</i> Miq. var. <i>siamensis</i>	-	50
<i>Sindora tonkinensis</i> K. Larsen & S.S. Larsen	+	72
<i>Smirnowia turkestanica</i> Bunge	+	150
<i>Smithia ciliata</i> Royle	+	76
<i>Sophora albescens</i> (Rehder) C.Y. Ma	+	72
<i>Sophora microphylla</i> Aiton	-	6
<i>Sophora mollis</i> (Royle) Baker ssp. <i>griffithii</i> (Stocks) Ali	+	9
<i>Sophora tetraptera</i> J.F. Mill.	+	89
<i>Spatholobus harmondii</i> Gagnep.	+	72
<i>Spatholobus parviflorus</i> (Roxb. ex DC.) Kuntze	+	76
<i>Stahlia monosperma</i> (Tul.) Urb.	-	68
<i>Stylosanthes angustifolia</i> Vogel	+	37
<i>Stylosanthes bracteata</i> Vogel	+	34
<i>Stylosanthes calcicola</i> Small	+	37
<i>Stylosanthes debilis</i> M.B. Ferreira & Sousa Costa	+	34
<i>Stylosanthes guianensis</i> (Aubl.) Sw. var. <i>intermedia</i> (Vogel) Hassl.	+	34
<i>Stylosanthes guianensis</i> (Aubl.) Sw. var. <i>marginata</i> Hassl.	+	34
<i>Stylosanthes guianensis</i> (Aubl.) Sw. var. <i>robusta</i> 't Mannetje	+	34
<i>Stylosanthes linearifolia</i> M.B. Ferreira & Sousa Costa	+	34
<i>Stylosanthes macrocarpa</i> S.F. Blake	+	37
<i>Stylosanthes macrocephala</i> M.B. Ferreira & Sousa Costa	+	34
<i>Stylosanthes pilosa</i> M.B. Ferreira & Sousa Costa	+	34
<i>Stylosanthes ruellioides</i> Mart. ex Benth.	+	34
<i>Stylosanthes sympodialis</i> Taub.	+	3
<i>Stylosanthes tomentosa</i> M.B. Ferreira & Sousa Costa	+	34
<i>Swainsona calcicola</i> Joy Thomps.	+	167
<i>Swainsona kingii</i> F. Muell.	+	167
<i>Swainsona leeana</i> J.Z. Weber	+	167
<i>Swainsona maccullochiana</i> F. Muell.	+	167
<i>Swainsona oroboides</i> F. Muell.	+	65
<i>Swainsona pterostylis</i> (DC.) Bakh. f.	+	167
<i>Swartzia apetala</i> Raddi	+	50
<i>Swartzia caribaea</i> Griseb.	+	142
<i>Swartzia jenmanii</i> Sandwith	+	150
<i>Swartzia pinnata</i> (Vahl) Willd.	-	108
<i>Swartzia simplex</i> (Sw.) Spreng. var. <i>ochracea</i> (DC.) R.S. Cowan	+	52
<i>Tachigali aurea</i> Tul.	+	69
<i>Tachigali guianensis</i> (Benth.) Zarruchi & Herend.	-	136
<i>Tachigali melinonii</i> (Harms) Zarucchi & Herend.	+	62
<i>Tachigali paniculata</i> Aubl. var. <i>alba</i> (Ducke) Dwyer	+	50
<i>Tachigali rugosa</i> (Mart. ex Benth.) Zarucchi & Pipoly	+	50
<i>Tadehagi triquetrum</i> (L.) H. Ohashi	+	38
<i>Tadehagi triquetrum</i> (L.) H. Ohashi ssp. <i>pseudotriquetrum</i> (DC.) H. Ohashi	+	72
<i>Tephrosia apollina</i> (Delile) Link	+	86

Taxon	Status ¹	Source ²
<i>Tephrosia baurei</i> Benth. ex A. Gray	+	65
<i>Tephrosia brachyodon</i> Domin	+	65
<i>Tephrosia bracteolata</i> Guill. & Perr.	+	42
<i>Tephrosia capensis</i> (Jacq.) Pers. var. <i>angustifolia</i> E. Mey.	+	61
<i>Tephrosia capensis</i> (Jacq.) Pers. var. <i>capensis</i>	+	61
<i>Tephrosia cephalantha</i> Baker	+	5
<i>Tephrosia dietrichii</i> Domin	+	65
<i>Tephrosia elongata</i> E. Mey. var. <i>tzaneenensis</i> (H.M. Forbes) Brummitt	+	60
<i>Tephrosia filipes</i> Benth.	+	65
<i>Tephrosia flagellaris</i> Domin	+	65
<i>Tephrosia juncea</i> R. Br. ex Benth.	+	38
<i>Tephrosia pallens</i> (Aiton) Pers.	+	61
<i>Tephrosia purpurea</i> (L.) Pers. ssp. <i>canescens</i> (E. Mey.) Brummitt	+	61
<i>Tephrosia purpurea</i> (L.) Pers. Var. <i>maritima</i> Haines	+	16
<i>Tephrosia rufula</i> Pedley	+	65
<i>Tephrosia supina</i> Domin	+	65
<i>Tephrosia virgata</i> H.M. Forbes	+	61
<i>Tephrosia zoutpansbergensis</i> Bremek.	+	61
<i>Tetraberlinia bifoliata</i> (Harms) Hauman	-	77
<i>Tetraberlinia moreliana</i> Aubr., v.	-	77
<i>Thermopsis barbata</i> Benth.	+	117
<i>Thermopsis lanceolata</i> R. Br.	+	150
<i>Trifolium africanum</i> Ser. var. <i>africanum</i>	+	153
<i>Trifolium africanum</i> Ser. var. <i>glabellum</i> (E. Mey.) Harv.	+	153
<i>Trifolium amabile</i> Kunth	+	146
<i>Trifolium canescens</i> Willd.	+	146
<i>Trifolium dasyphyllum</i> Torr. & A. Gray	+	19
<i>Trifolium decorum</i> Chiov.	+	55
<i>Trifolium echinatum</i> M. Bieb. Var. <i>carmeli</i> (Boiss.) Gibelli & Belli	+	9
<i>Trifolium haydenii</i> Porter	+	5
<i>Trifolium heldreichianum</i> (Gibelli & Belli) Hausskn.	+	146
<i>Trifolium lanceolatum</i> (J.B. Gillett) J.B. Gillett	+	27
<i>Trifolium mattirolanum</i> Chiov.	+	150
<i>Trifolium nanum</i> Torr.	+	92
<i>Trifolium parryi</i> A. Gray	+	92
<i>Trifolium plebeium</i> Boiss.	+	150
<i>Trifolium procumbens</i> L.	+	5
<i>Trifolium quartinianum</i> A. Rich.	+	55
<i>Trifolium scutatum</i> Boiss.	+	150
<i>Trifolium simense</i> Fresen.	+	38
<i>Trigonella cretica</i> (L.) Brot.	-	60
<i>Trigonella glabra</i> Thunb. subsp. <i>uncata</i> (Boiss. & Noe) Lassen	+	9
<i>Uraria cylindracea</i> Benth.	+	65
<i>Uraria lagopus</i> DC.	+	76
<i>Uraria lagopus</i> DC. var. <i>neglecta</i> (Prain) H. Ohashi	+	76
<i>Uraria rufescens</i> (DC.) Schindl.	+	76
<i>Vandasina retusa</i> (Sol. ex Benth.) Rauschert	+	150
<i>Vatairea heteroptera</i> (Allemão) Ducke	-	50
<i>Vicia amurensis</i> Oett.	+	150
<i>Vicia bungei</i> Ohwi	+	169
<i>Vicia faba</i> L. var. <i>major</i> (Harz.) Beck	+	168
<i>Vicia faba</i> L. var. <i>minuta</i> (Hort. ex Alef.) Mansf.	+	168
<i>Vicia pseudoorobus</i> Fisch. & C.A. Mey.	+	150
<i>Vicia ramuliflora</i> (Maxim.) Ohwi	+	150
<i>Vicia serratifolia</i> Jacq.	+	146
<i>Vicia sylvatica</i> L.	+	150
<i>Vicia venosa</i> (Willd. ex Link) Maxim.	+	150
<i>Vigna comosa</i> Baker ssp. <i>comosa</i>	+	49

Taxon	Status ¹	Source ²
<i>Vigna cylindrica</i> (L.) Skeels	+	146
<i>Vigna multinervis</i> Hutch. & Dalziel	+	49
<i>Vigna nakashimae</i> (Ohwi) Ohwi & O. Ohashi	+	123
<i>Vigna oblongifolia</i> A. Rich.	+	38
<i>Vigna radiata</i> (L.) R. Wilczek var. <i>radiata</i>	+	5
<i>Vigna unguiculata</i> (L.) Walp. ssp. <i>unguiculata</i>	+	39
<i>Wiborgia incurvata</i> E. Mey.	+	61
<i>Wiborgia monoptera</i> E. Mey.	+	60
<i>Zapoteca tetragona</i> (Willd.) H.M. Hern.	-	150
<i>Zenkerella citrina</i> Taub.	-	150
<i>Zornia brasiliensis</i> Vogel	+	38
<i>Zygia inaequalis</i> (Willd.) Pittier	+	50
<i>Zygia juruana</i> (Harms) L. Rico	+	150

¹ Status: +, root nodules reported as present; -, root nodules reported as absent.

² Source:

1. Abdelmounen, H., Filali-Maltouf, A., Neyra, M., Belabed, A., and Missbah El Idrissi, M. 1999. Effect of high salts concentrations on the growth of rhizobia and responders to added osmotica. *Journal of applied Microbiology* 86:889-898.
2. Aguirre, M.J., M. Valdés, and R.M. Silvestre-Bradley. 1988. Simbiosis entre rizobios y cuatro leguminosas tropicales adaptadas en Chiapas, México. *Pasturas tropicales* 10(3): 18-21.
3. Ahmed, B., and J.M. Keoghan. 1978. Nitrogen fixation by some forage legumes under West Indian conditions. In J. Dobereiner, R.H. Burris, R.H. Hollaender, A.A. Franco, C.A. Neyra, and D.B. Scott, eds., *Limitations and potentials for biological nitrogen fixation in the tropics*. Basic life Sciences 10: 342-343. [Abstracts of original papers]
4. Alazard, D. 1985. Stem and root nodulation in *Aeschynomene* spp. *Applied and Environmental Microbiology* 50(3): 732-734.
5. Allen, O.N., and E. Hamatová. 1973. *IBP World Catalogue of Rhizobium Collections*. F.A. Skinner, ed. 282 pp. International Biological Programme, London.
6. Aronson, J., C. Ovalle, and J. Avendaño. 1992. Early growth rate and nitrogen fixation potential in forty-four legume species grown in an acid and a neutral soil from central Chile. *Forest Ecology and Management* 47: 225-244.
7. Assefa, F., and D. Kleiner. 1998. Nodulation pattern and acetylene reduction (Nitrogen fixation) activity of some highland and lowland *Acacia* species of Ethiopia. *Biol. Fertil. Soils* 27: 60-64.
8. Assefa, F., and Kleiner, D. 1997. Nodulation of African yam bean (*Sphenostylis stenocarpa*) by *Bradyrhizobium* sp. isolated from *Erythrina brucei*. *Biology and Fertility of Soils* 25:209-210.
9. Athar, M., and A. Mahmood. 1985. Qualitative study of the nodulating ability of legumes of Paskistan: list 3. *Trop. Agric.* 62(1): 49-51.
10. Baggie, I., Zapata, F., Sanginga, N., and Danso, S.K.A. 2000. Ameliorating acid infertile rice soil with organic residue from nitrogen fixing trees. *Nutrient Cycling in Agroecosystems* 57:183-190.
11. Ballard, R.A., and A. Charman. 2000. Nodulation and growth of pasture legumes with naturalised soil rhizobia – 1. *Annual Medicago* spp. *Australian Journal of Experimental Agriculture* 40: 939-948.
12. Barrera, L.L., Trujillo, M.E., Goodfellow, M., García, F.J., Hernández-Lucas, I., Dávila, G., Berkum, P. van, and Martínez-Romero, E. 1997. Biodiversity of bradyrhizobia nodulating *Lupinus* spp. *International Journal of Systematic Bacteriology* 47:1086-1091.
13. Basak, M.K., and S.K. Goyal. 1980. Studies on the biology of tree legumes – *Rhizobium* symbiosis: Nodulation pattern and cross inoculation trials with the tree legumes and cultivated legumes. *Annals of Arid Zone* 19(4): 427-431.
14. Basak, M.K., and S.K. Goyal. 1980. Studies on tree legumes. II. Further additions to the list of nodulating tree legumes. *Plant and Soil* 56: 33-37.
15. Béreau, M., and Garbaye, J. 1994. First observations on the root morphology and symbioses of 21 major tree species in the primary tropical forest rain forest of French Guyana. *Annales des Sciences Forestiere* 51:407-416.
16. Biswal, A.K., L.M. Sahoo, A.K. Mishra, and B. Padhi. 1987. Nodulation, total nitrogen and crude protein content of some leguminous plants of different ecological conditions of Orissa. *Environment and Ecology* 5(4): 694-699.
17. Bowen, D., Dart, P.J., Ryan, P., and Harwood, C.E. 1998. *Rhizobium* inoculants for Australian *Acacia* species. In J.W. Turnbull, H.R. Crompton, and K. Pinyopusarerk, eds., *Recent Developments in Acacia Planting: Proceedings of an international workshop held in Hanoi, Vietnam, 27-30 October 1997*, pp. 368-370. Australian Centre for International Agricultural Research, Canberra, Australia.
18. Bowen, E.J. 1981. The effect of nitrogen fertilizer on the establishment and vigour of Oxley fine-stem stylo. *Tropical Grasslands* 15(3): 140-144.
19. Bowman, W.D., Schardt, J.C., and Schmidt, S.K. 1996. Symbiotic N₂-fixation in alpine tundra: ecosystem input and variation in fixation rates among communities. *Oecologia* 108:345-350.
20. Brewbaker, J.L. 1987. Significant nitrogen fixing trees in agroforestry systems. *Agroforestry: realities, possibilities and potentials*, pp. 31-45. Martinus Nijhoff, Dordrecht, Netherlands.

21. Brockwell, J. 1998. Match rhizobia and temperate species of *Acacia*. In J.W. Turnbull, H.R. Crompton, and K. Pinyopusarerk, eds., Recent Developments in *Acacia* Planting: Proceedings of an international workshop held in Hanoi, Vietnam, 27-30 October 1997, pp. 264-273. Australian Centre for International Agricultural Research, Canberra, Australia.
22. Brockwell, J., Bottomley, P.J., and Thies, J.E. 1995. Manipulation of Rhizobia microflora for improving legume productivity and soil fertility: A critical assessment. *Plant and Soil* 174:143-180.
23. Brown, S.M., and Walsh, K.B. 1994. Anatomy of the legume nodule cortex with respect to nodule permeability. *Australian Journal of Plant Physiology* 21:49-68.
24. Brown, S.M., and Walsh, K.B. 1994. Anatomy of the legume nodule cortex with respect to nodule permeability. *Australian Journal of Plant Physiology* 21:49-68.
25. Burdon, J.J., Gibson, A.H., Searle, S.D., Woods, M.J., and Brockwell, J. 1999. Variation in the effectiveness of symbiotic associations between native rhizobia and temperate Australian *Acacia*: within-species interactions. *Journal of Applied Ecology* 36:398-408.
26. Burton, J.C. 1977. Associations of some uncommon leguminous plants with various rhizobia. In J.M. Vincent, A.S. Whitney, and J. Bose, eds., Exploiting the legume *Rhizobium* symbiosis in tropical agriculture, pp. 381-384. Hawaii University, Hawaii.
27. Burton, J.C. 1979. New developments in inoculating legumes. 1979. In Subba Rao, N.S., ed., Recent advances in Biological Nitrogen Fixation, pp. 380-405. Oxford & IBH Publishing Co., New Delhi, India.
28. Bushby, H.V.A., Date, R.A., Norris, D.O., and Panitz, R.B. 1984. *Rhizobium* screening for a range of crop and pasture legumes: lists of strains of *Rhizobium* and host legumes forming effective nitrogen fixing associations. 7:1-39.
29. Carr, D.J., S.G.M. Carr, and W.R. Papst. 1980. Field studies of nitrogen fixation of Australian alpine plants and soils. *Australian Journal of Ecology* 5: 211-220.
30. Chada, S. de S., and H. de Polli. 1988. Nodulação de leguminosas tropicais proissoras para adubação verde em solo deficiente em fósforo. *Pesq. Agropec. Bras.* 23(11): 1197-1202.
31. Chaukiyal, S.P., Himmat Singh, K.C., and Pokhriyal, T.C. 1999. Effects of seasonal variations on nodulation and nitrogen fixation behaviour in some *Acacia* species. *Annals of Forestry* 7(1):112-119.
32. Chen, W.X., Wang, E.T., Wang, S.Y., Li, Y.B., Chen, X.Q., and Li, Y. 1996. Characteristics of *Rhizobium tianshanense* sp. nov. a moderate- and slow-growing root nodule bacterium isolated from arid-saline environment in Xinjiang, China. In F.D. Li, T.A. Lie, W.X. Chen, and J.C. Zhou, eds., Diversity and Taxonomy of Rhizobia, pp. 49-58. China Agricultural Sciencetech Press, Beijing, China.
33. Cocks, M.P., and Stock, W.D. 2001. Field patterns of nodulation in fifteen *Aspalathus* species and their ecological role in the fynbos vegetation of southern Africa. *Basic and Applied Ecology* 2:115-125.
34. Costa, N.M.S. 1981. Native legumes in Minas Gerais state, Brazil. In P.H. Graham and S.C. Harris, eds., Biological nitrogen fixation technology for tropiucal agriculture, pp. 337-346. Centro Internacioanl de Agricultura Tropical, Cali, Colombia.
35. Craig, G.F., D.T. Bell, and C.A. Atkins. 1990. Response to slat and waterlogging stress of ten taxa of *Acacia* selected from naturally saline areas of Western Australia. *Aust. J. Bot.* 38: 619-630.
36. Dart, P., M. Umali-Garcia, and A. Alemdras. 1991. Role of symbiotic associations in nutrition of tropical acacias. In J.W. Turnbull, ed., Advances in tropical acacia research. Proceedings of an international workshop held in Bangkok, Thailand, 11-15 February 1991, pp. 13-19. Australian Centre for International Agricultural Research, Canberra, Australia.
37. Date, R.A., and D.O. Norris. 1979. *Rhizobium* screening of *Stylosanthes* species for effectiveness in nitrogen fixation. *Aust. J. Agric. Res.* 30: 85-104.
38. Date, R.A., Bushby, H.V.A., and Panitz, R.B. 1984. The CB *Rhizobium* strain catalogue. *Genetic Resources Communication* 5:1-33.
39. Dayap, F.T., and E. T. Rasco, Jr. 1988. Genetic analysis for nitrogen fixation (acetylene reduction) and nodulation in bush sitao (*Vigna unquiculata* ssp. *sesquipedalis* (L.) Verdc. × *Vigna unquiculata* (L.) Walp. ssp. *unquiculata*). *Philippine Journal of Crop Science* 13: Supplement 1, S18.
40. Deschodt, C.C., and Strijdom, B.W. 1976. Effective noldulation of *Aspalathus linearis* ssp. *linearis* by rhizobia from other *Aspalathus* species. *Phytophylactica* 8:103-104.
41. Ding, M.-M., Wang, Z.-M., Yi, W.-M., and Yu, Z.Y. 1994. Characteristics and effectiveness of tree legume rhizobia. *Nitrogen Fixing Tree Research Reports* 12:18-20.
42. Doignon-Bourcier, F., Sy, A., Willems, A., Torck, U., Dreyfus, B., Gillis, M., and Lajudie, P. de. 1999. Diversity of *Badyrhizobia* from 27 tropical Leguminosae species native of Senegal. *System. Appl. Microbiol.* 22: 647-661.
43. Domingo, I.L. 1983. Nitrogen fixation in Southeast Asian forestry; research and practice. In J.C. Gordon and C.T. Wheeler, eds., Biological nitrogen fixation in forest ecosystems: foundations and applications, pp. 295-315. Martinus Nijhoff/Dr. W. Junk Publishers, The Hague.
44. Dommergues, Y., Duhoux, E., and Diem, H.G. 1999. Les arbres fixateurs d'azote. Editions Espaces, Montpellier, France.
45. Dreyfus, B.L., and Y.R. Dommergues. 1981. Nodulation of *Acacia* species by fast- and slow-growing tropical strains of *Rhizobium*. *Applied and Environmental Microbiology* 41(1): 97-99.

46. Eaglesham, A.R.J., and A.A. Szalay. 1983. Aerial stem nodules on *Aeschynomene* spp. *Plant Science Letters* 29: 265–272.
47. Elkan, G.H., J.C. Wynne, and T.J. Schneeweis. 1981. Isolation and evaluation of strains of *Rhizobium* collected from centres of diversity in South America. *Trop. Agric. (Trinidad)* 58(4): 297–305.
48. Eskew, D.L., and I.P. Ting. 1978. Nitrogen fixation by legumes and blue-green algal-lichen crusts in a Colorado desert environment. *Amer. J. Bot.* 65(8): 850–856.
49. Ezedinma, F.O.C., N.N. Agnim, and S.S.C. Onyekwelu. 1979. The contribution of symbiotic nitrogen fixation to the nitrogen economy of natural ecosystem. I. Occurrence of herbaceous legumes in derived savanna fallow and their nodulation in pot culture. *Plant and Soil* 51: 503–513.
50. Faria, S.M. de, H.C. de Lima, A.A. Franco, E.S.F. Mucci, and J.I. Sprent. 1987. Nodulation of legume trees from South East Brazil. *Plant and Soil* 99: 347–356.
51. Faria, S.M. de, Lima, H.C. de, and Olivares, F.L. 1999. Nodulação em espécies florestais especificidade hospedeira e implicações na sistemática de Leguminosae. In J.O. Siqueira et al., eds., *Soil fertility, soil biology and plant nutrition interrelationships*, pp. 667–686. Sociedade Brasileira de Ciência do Solo, Universidade Federal da Laurus Brazil, Viçosa, Brazil.
52. Faria, S.M. de, McInroy, S.G., and Sprent, J.I. 1987. The occurrence of infected cells, with persistent infection threads, in legume root nodules. *Canadian Journal of Botany* 65:553–558.
53. Faria, S.M. de, Sutherland, J.M., and Sprent, J.I. 1986. A new type of infected in root nodules of *Andira* spp. (Legumonisaeae). *Plant Science* 45:143–147.
54. Felker, P., and P.R. Clark. 1980. Nitrogen fixation (acetylene reduction) and cross inoculation in 12 *Prosopis* (mesquite) species. *Plant and Soil* 57: 177–186.
55. Fredericks, J.B. 1990. Evaluation of African *Trifolium* species for growth and biological nitrogen fixation. *Dissertation Abstracts International* 50(7): 2682-B.
56. Frioni, L., Malat,s, D., Irigoyen, I., and Doderá. 1998. Promiscuity for nodulation and effectivity in the N₂-fixing legume tree *Acacia caven* in Uruguay. *Applied Soil Ecology* 7:239–244.
57. Gao, J.L., Sun, J.G., Li, Y., Wang, E.T., and Chen, W.X. 1994. Numerical taxonomyh and DNA relatedness of tropical rhizobia isolated from Hainan Province, China. *International Journal of Systematic Bacteriology* 44(1):151–158.
58. Gault, R.R., Pilka, A., Hedd, D.M., and Brockwell, J. 1994. Nodulation studies on legumes exotic to Australia: symbiotic relationships between *Chamaecytisus plamensis* (tagasaste) and *Lotus* spp. *Australian Journal of Experimental Agriculture* 34:385–394.
59. Gibson, A.C., Sharifi, M.R., and Rundel, P.W. 1998. Ecophysiological observations on Lane Mountain milkvetch, *Astragalus jargerianus* (Fabaceae), a proposed endangered species of the Mojave Desert. *Aliso* 17(1):77–82.
60. Grobbelaar, N., and M.W. van Rooyen. 1979. A qualitative study of the nodulating ability of legume species: List 5. *J. S. African Bot.* 45(3): 267–272.
61. Grobbelaar, N., M.W. van Rooyen, and N. van Rooyen. 1983. A qualitative study of the nodulating ability of legume species: List 6. *S. Afr. J. Bot.* 2: 329–332.
62. Guehl, J.M., Domenach, A.M., Bereau, M., Barigah, T.S., Casabianca, H., Ferhi, A., and Garbaye, J. 1998. Functional diversity ijn an Amazonian rainforest of French Guyana: a dual isotope approach (delta15N and delta13C). *Oecologia* 116:316–330.
63. Guinto, D.F., Xu, Z.-H., House, A.P.N., and Saffigna, P.G. 2000. Assessment of N₂ fixation by understory acacias in recurrent burnt eucalypt forests of subtropical Australia using 15N isotope dilution techniques. *Canadian Journal of Forest Research* 30:112–121.
64. Gwaze, D.P., H.T.L. Stewart, and R.A. Snieszko. 1988. Variation in height growth and nodulation of *Acacia albida* and *A. erioloba* seedlings from Zimbabwe seed sources. *Nitrogen Fixing Tree Research Reports* 6: 34–36.
65. Hacker, J.B. 1990. A guide to herbaceous and shrub legumes of Queensland. University of Queensland Press, St. Lucia, Australia.
66. Hadri, A.-E., Spaink, H.P., Bisseling, T., and Brewin, N.J. 1998. Diversity of root nodulation and rhizobial infection processes. In H.P. Spaink, A. Kondorosi, and P.J.J. Hooykass, eds., *The Rhizobiaceae: Molecular biology of model plant-associated bacteria*, pp. 347–360. Kluwer Academic Publishers,
67. Hailu, T., Negash, L., and Olsson, M. 2000. *Millettia ferruginea* from southern Ethiopia: Impasts on fertility and growth of maize. *Agroforestry Systems* 48:9–24.
68. Halliday, J. 1984. Register of nodulation reports for leguminous trees and other arboreal genera with nitrogen fixing members. *Nitrogen Fixing Tree Research Reports* 2: 38–45.
69. Halliday, J., and Nakao, P.L. 1982. Master list of woody species under consideration as nitrogen fixing trees. In J.L. Brewbaker, ed., *Resource Documents on Nitrogen-Fixing Trees*, NFTA 82-03/12, pp. 4–11. Nitrogen Fixing Tree Association, Waimanalo, Hawaii.
70. Halliday, J., and P. Somasegaran. 1983. Nodulation, nitrogen fixation, and *Rhizobium* strain affinities in the genus *Leucaena*. *Leucaena research in the Asian Pacific region*, pp. 27–32. International Development Research Centre, Ottawa, Canada.
71. Han, S.-F. 1995. A nodulation survey of tree legumes in China. *Journal of Nanjing Forestry University* 19(4):89–92.

72. Han, S.-F., and Zhou, X.-Q. 1990. Register of nodulation reports for leguminous trees and shrubs in China. *Journal of Nanjing Forestry University* 14(3):84-90.
73. Han, S.-F., and Zhou, X.-Q. 1992. Studies on the possibility to induce nodulation and nitrogen fixation of non-nodulated tree legumes. *Journal of Nanjing Forestry University* 16(1):19-24.
74. Harrier, L.A., P. Whitty, J.M. Sutherland, and J.I. Sprent. 2000. Pre-infection events in non-nodulating species of African *Acacia*. *Afr. Jr. Ecol.* 38: 8–15.
75. Hashem, F. 2001. Pers. comm.
76. Himmat Singh, K.C., and Pokhriyal, T.C. 1997. Survey and identification of some leguminous nitrogen fixing plants in eastern part of Nepal. *Annals of Forestry* 5(2):205-213.
77. Hogberg, P., and Alexander, I.J. 1995. Roles of root symbioses in African woodland and forest: evidence from ^{15}N abundance and foliar analysis. *Journal of Ecology* 83:217-224.
78. Högborg, P., and J.-E. Nylund. 1981. Ectomycorrhizae in coastal Miombo woodland of Tanzania. *Plant and Soil* 63: 283–289.
79. Hogberg, P., and Pearce, G.D. 1986. Mycorrhizas in Zambian trees in relation to host taxonomy, vegetation type and successional patterns. *Journal of Ecology* 74:775-785.
80. Holton, B. Jr., M.G. Barbour, and S.N. Martens. 1991. Some aspects of the nitrogen cycle in a Californian strand ecosystem. *Madroño* 38(3): 170–184.
81. Howieson, J.G., Fillery, I.R.P., Legocki, A.B., Sikorski, M.M., Stepkowski, T., Minchin, F.R., and Dilworth, M.J. 1998. Nodulation, nitrogen fixation and nitrogen balance. In J.S. Gladstones, C.A. Atkins, and J. Hamblin, eds., *Lupins as crop plants: Biology, production and utilization*, pp. 149-180. CAB International, Wallingford, United Kingdom.
82. Idrissi, M.M. El, Aujjar, N., Belabed, A., Dessaux, Y., and Filali-Maltouf, A. 1996. Characterization of rhizobia isolated from Carob tree (*Ceratonia siliqua*). *Journal of Applied Bacteriology* 80:165-173.
83. Izaguirre-Mayoral, M.L., O. Carballo, S. Flores, M.S. de Mallorca, and T. Oropeza. 1992. Quantitative analysis of the symbiotic N_2 -fixation, non-structural carbohydrates and chlorophyll content in sixteen native legume species collected in different savanna sites. *Symbiosis* 12: 293–312.
84. Izaguirre-Mayoral, M.L., Vivas, A.I., and Oropeza, T. 1995. New insights into the symbiotic performance of native tropical legumes: I. Analysis of the response of thirty-seven native legume species to artificial shade in a neotropical savanna. *Symbiosis* 19:111-129.
85. Jacot, K.A., Lüscher, A., Nüssberger, J., and Hartwig, U.A. 2000. Symbiotic N_2 fixation of various legume species along an altitudinal gradient in the Swiss Alps. *Soil Biology & Chemistry* 32(8/9):1043-1052.
86. Jain, P.S., and P. Shrivastava. 1984. Development and structure of ineffective nodules in some leguminous weeds. *Proc. Indian Acad. Sci. (Plant Sci.)* 93(4): 467–470.
87. James, E.K., Iannetta, P.P.M., Naisbitt, T., Goi, S.R., Sutherland, J.M., Sprent, J.I., Minchin, F.R., and Brewin, N.J. 1994. A survey of N_2 -fixing nodules in the Leguminosae with particular reference to intercellular glycoproteins and the control of oxygen diffusion. *Proceedings of the Royal Society of Edinburgh* 102B:429-432.
88. James, E.K., Loureiro, M. de F., Pott, A., Pott, V.J., Martins, C.M., Franco, A.A., and Sprent, J.I. 2001. Flooding-tolerant legume symbioses from the Brazilian Pantanal. *New Phytologist* 150(3):723-739.
89. Jarvis, B.D.W., Pankhurst, C.E., and Patel, J.J. 1982. *Rhizobium loti*, a new species of legume root nodule bacteria. *International Journal of Systematic Bacteriology* 32(3):378-380.
90. Jiang, R.W., Zhou, R., Jiang, M.L., and Zhang, X.J. 1988. Evaluation of the symbiotic potential of some wild *Arachis* species with two *Rhizobium* strains. *International Arachis Newsletter* 4: 23–24.
91. Johnson, D.A., and M.D. Rumbaugh. 1981. Nodulation and acetylene reduction by certain Rangeland legume species under field conditions. *Journal of Rangeland Management* 34(3): 178–181.
92. Johnson, D.A., and M.D. Rumbaugh. 1986. Field nodulation and acetylene reduction activity of high-altitude legumes in the western United States. *Arctic and Alpine Research* 18(2): 171–179.
93. Khasa, P., V. Furlan, and K. Lumande. 1990. Symbioses racinaires chez quelques essences forestières importantes au Zaïre. *Bois et Forêts des Tropiques* 224: 27–33.
94. Khbaya, B., Neyra, M., Normand, P., Zerhari, K., Filali-Maltouf, A. 1998. Genetic diversity and phylogeny of rhizobia that modulate *Acacia* spp. in Morocco assessed by analysis of rRNA genes. *Applied and Environmental Microbiology* 64(12):4912-4917.
95. Kishinevsky, B.D., and Kishinevsky, S. 1996. Characterization of root-nodule bacteria isolated from various *Hedysarum* species. In F.D. Li, T.A. Lie, W.X. Chen, and J.C. Zhou, eds., *Diversity and Taxonomy of Rhizobia*, pp. 49-58. China Agricultural Sciencetech Press, Beijing, China.
96. Kjaer, S. 1992. Biological nitrogen fixation in *Pachyrhizus ahipa* (Wedd.) Parodi. *Annals of Botany* 70: 11–17.
97. Kothandaraman, R., Premakumari, D., and Sivasankara Panickar, P.K. 1987. Studies on growth, nodulation and nitrogen fixation by *Mucuna bracteata* D.C. In R. Kothandaram, J. Mathew, A.K. Krishnakumar, J. Kochuthrassiamma, K. Jayarathnam, and M.J. Seturaj, eds., *Proceedings of the sixth Symposium on plantation crops*, pp. 283-288. Kottayam, India.
98. Ladha, J.K., Pareek, R.P., and Becker, M. 1992. Stem-nodulating legume-*Rhizobium* symbiosis and its agronomic use in lowland rice. *Advances in Soil Science* 20:147-192.

99. Ladipo, D.O. 1994. *Dalbergia* species of West Africa: botany, distribution and potential for agroforestry. In S.B. Westley and J.M. Rossetko, eds., *Dalbergia*: Proceedings of an international workshop, pp. 27-37. Winrock International, Morrilton.
100. Lafay, B., and J.J. Burdon. 2001. Small-subunit rDNA genotyping of *Rhizobia* nodulating Australian *Acacia* spp. *Applied and Environmental Microbiology* 67(1): 396-402.
101. Laguerre, G., Berkum, P. van, Amarger, N., and Prévost, D. 1997. Genetic diversity of rhizobial symbionts isolated from legume species within the genera *Astragalus*, *Oxytropis*, and *Onobrychis*. *Applied Environmental Microbiology* 63:4748-4758.
102. Lajudie, P. de, Willems, A., Nick, G., Mohamed, S.H., Torck, U., Coopman, R., Filali-Maltouf, A., Kersters, K., Creyfus, B., Linström, K., and Gillis, M. 1999. *Agrobacterium* bv. 1 strains isolated from nodules of tropical legumes. *Systematic and Applied Microbiology* 22:119-132.
103. Lajudie, P. de, Willems, A., Nick, G., Moreira, F., Molouba, F., Hoste, B., Torck, U., Neyra, M., Collins, D., Lindström, K., Dreyfus, B., and Gillis, M. 1998. Characterization of tropical tree rhizobia and description of *Mesorhizobium plurifarium* sp. nov. *International Journal of Systematic Bacteriology* 48:369-382.
104. Lamont, B. 1982. Specialised roots in the genus *Acacia*: A review. *Annual Report Mulga Research Centre* 5: 9-11.
105. Langkamp, P.J., L.B. Swinden, and M.J. Dalling. 1982. Nutrient fixation (acetylene reduction) by *Acacia pellita* on areas restored after mining at Groote Eylandt, Northern Territory. *Austral. J. Bot.* 27: 353-361.
106. Lawrie, A.C. 1981. Nitrogen fixation by native Australian legumes. *Austral. J. Bot.* 29: 143-157.
107. Lim, G. 1979. *Rhizobium* and nodulation of legumes in Singapore. In W.J. Broughton, C.K. John, J.C. Rajaro, and B. Lim, eds., *Soil microbiology and plant nutrition*, pp. 159-175. University of Malaya Press, Kuala Lumpur, Malaya.
108. Lim, G. 1983. *Rhizobia* and nitrogen-fixing trees in Singapore. In IDRC, Ottama, Canada, ed., *Laeucaena* Research in the Asian - Pacific Region: proceedings of a workshop held in Singapore, 23-26 Nov. 1982, pp. 33-37. IDRC, Ottawa, Canada.
109. Loureiro, M. de F., Faria, S.M. de, Baldani, J.I., and Franco, A.A. 1993. Morphological and symbiotic characteristics of stem nodules of *Discolobium* spp. and its rhizobia. In P. Palacios, J. Mora, and W.E. Newton, eds., *New horizons in nitrogen fixation*, p. 628. Kluwer Academic Publishers, Dordrecht, The Netherlands.
110. Luyindula, N., and Haque, I. 1994. Early nodulation, growth, and nitrogen fixation in *Sesbania* species as affected by phosphorus, nitrogen, and rhizobia inoculation in a vertisol. *Nitrogen Fixing Tree Research Reports* 12:134-138.
111. Mahmood, A. 1999. A comparison of nitrogen concentration between wild and cultivated legumes of Sindh. *Pakistan Journal of Botany* 31(1): 183-192.
112. Manninger, E., I. Csapody, and E. Bakondi-Zámory. 1984. Wild wachsende Leguminosen und das Vorkommen der durch *Rhizobium*-Bakterien gebildeten Wurzelknöllchen in Wald- und Wiesenökosystemen. *Zbl. Mikrobiol.* 139: 319-326.
113. Martínez-Romero, E., and Caballero-Mellado, J. 1996. *Rhizobium* phylogenies and bacterial genetic diversity. *Critical Reviews in Plant Sciences* 15(2): 113-140.
114. McColl, J.G., and R.L. Edmonds. 1983. Symbiotic nitrogen fixation by *Daviesia mimsoides* under *Eucalyptus*. General Technical Report, Pacific Northwest Forest and Range Experiment Station, USDA Forest Service, no. 163, pp. 122-129.
115. Michiels, J., Dombrecht, B., Vermeiren, N., Xi, C.-W., Luyten, E., and Vanderleyden, J. 1998. *Phaseolus vulgaris* is a non-selective host for nodulation. *FEMS Microbiology Ecology* 26:193-205.
116. Mullen, B.F., Frank, V.E., and Date, R.A. 1998. Specificity of rhizobial strains for effective N₂ fixation in the genus *Leucaena*. *Tropical Grasslands* 32:110-117.
117. Nair, S., P.K. Jha, M.C. Gopinathan, and C.R. Babu. 1988. New nodulating papilionoid legumes of potential value for revegetation of degraded ecosystems. *Proc. Indian Acad. Sci. (Plant Sci.)* 98(1): 41-48.
118. Newton, W.E., and W.H. Orme-Johnson. 1980. Nitrogen fixation, volume 11: Symbiotic associations and Cyanobacteria. University park press, Baltimore.
119. Nick, G., Lajudie, P. de, Eardly, B.D., Suomalainen, S., Paulin, L., Zhang, X.-P., Gillis, M., and Lindström, K. 1999. *Sinorhizobium arboris* sp. nov. and *Sinorhizobium kostiense* sp. nov., isolated from leguminous trees in Sudan and Kenya. *International Journal of Systematic Bacteriology* 49:1359-1368.
120. Novikova, N.I., Pavlova, E.A., Vorobjev, N.I., and Linesshchenko, E.V. 1994. Numerical taxonomy of *Rhizobium* strains from legumes of the temperate zone. *International Journal of Systematic Bacteriology* 44(4): 734-742.
121. Odee, D.W., Sutherland, J.M., Kimiti, J.M., and Sprent, J.I. 1995. Natural rhizobial populations and nodulation status of woody legumes growing in diverse Kenyan conditions. *Plant and Soil* 173:211-224.
122. Ogan, M.T. 1990. The nodulation and nitrogenase activity of natural stands of mangrove legumes in a Nigerian swamp. *Plant and Soil* 123: 125-129.
123. Oh, H.-S., Chun, J.-Y., Lee, M.-S., Min, K.-H., Lee, S.-H., and Cheon, C.-I. 2000. Role of *hsfA* gene on host-specificity by *Bradyrhizobium japonicum* in a broad range of tropical legumes. *Canadian Journal of Microbiology* 46:81-84.
124. Ovalle, C., Aronson, J., Pozo, A. del, and Avendao, J. 1999. Restoration and rehabilitation of mixed espinales in central Chile: 10-year report and appraisal. *Arid Soil Research and Rehabilitation* 13:369-381.
125. Pankhurst, C.E., and Jones, W.T. 1979. Effectiveness of *Lotus* root nodules. 11. Relationship between root nodule effectiveness and 'in vitro' sensitivity of fast-growing *Lotus* rhizobia to flavolans. *Journal of Experimental Botany* 30(119): 1095-1107.

126. Parker, M.A., and A. Link. 2000. Relationships of bradyrhizobia from *Platypodium* and *Machaerium* (Papilionoideae: tribe Dalbergieae) on Barro Colorado Island, Panama. *International Journal of Systematic and Evolutionary Microbiology* 50: 1179–1186.
127. Pärsim, E. 1967. Acid production by fermentation of carbohydrates by *Rhizobium*. *Izy. Akad. Nauk. Estonsk. SSR XVI Biol.* 2: 163–174. [In Russian with English summary.]
128. People, M.B., D.F. Herridge, and F.J. Bergersen. 1988. Measurement of nitrogen fixation in crop and shrub legumes. *In Green manure in rice farming*, pp. 223–237. International Rice Research Institute, Manila.
129. Peoples, M.B., Atkins, C.A., Pate, J.S., Chong, K., Faizah, A.W., Suratmini, P., Nurhayati, D.P., Bagnall, D.J., and Bergersen, F.J. 1991. Re-evaluation of the role of ureides in the xylem transport of nitrogen in *Arachis* species. *Physiologia Plantarum* 83:560–567.
130. Pokhriyal, T.C., Chaukiyal, S.P., and Naithani, H.B. 1997. Nodule biomass and noitrogen fixation studies in some leguminous nitrogen fixing plants. *Indian Forester* 123:1197–1198.
131. Pokhriyal, T.C., Chaukiyal, S.P., Naithani, H.B., and Gupta, B.B. 1992. Identification of some nitrogen fixing herb and shrub species from Dehra Dun-Mussoorie region. *Indian Forester* 118:681–683.
132. Prameela, K., and Charya, M.A.S. 1999. Impact of coal mining on the growth of certain *Cassia* species. *Geobios* 26(4):199–204.
133. Prévost, D., L.M. Bordeleau, S. Caudry-Reznick, H.M. Schulman, and H. Antoun. 1987. Characteristics of rhizobia isolated from three legumes indigenous to the Canadian high artic: *Astragalus alpinus*, *Oxytropis maydelliana*, and *Oxytropis arctobia*. *Plant and Soil* 98: 313–324.
134. Radomiljac, A.M., and McComb, J.A. 1998. Nitrogen-fixing and non-nitrogen-fixing woody host influences on the growth of the root hemi-parasite *Santalum album* L. *In Proceedings of an international seminar held on 18-19 December 1997 at the Institute of Wood Science and Technology, Bangalore, India*, pp. 54–57. Australian Centre for International Agricultural Research, Canberra, Australia.
135. Reddell, P., and A.R. Milnes. 1992. Mycorrhizas and other specialised nutrient-acquisition strategies: their occurrence in woodland plants from Kakadu and their role in rehabilitation of waste rock dumps at a local uranium mine. *Austral. J. Bot.* 40: 223–242.
136. Roggy, J.C., Pr, vost, M.F., Gourbiere, F., Casabianca, H., Garbaye, J., and Domenach, A.M. 1999. Leaf natural ¹⁵N abundance and total N concentration as potential indicators of plant N nutrition in legumes and pioneer species in a rain forest of French Guiana. *Oecologia* 120:171–182.
137. Roshetko, J.M., Dagar, J.C., Puri, S., Khandale, D.Y., Takawale, P.S., Bheemaiah, G., and Basak, C. 1996. Selecting species of nitrogen fixing fodder trees. *In O.I. Rosketko and R.C. Gutteridge, eds., Nitrogen fixing trees for fodder production: a field manual*, pp. 19–24. Forest, Farm, and Community Tree Network, Morrilton, Arkansas.
138. Roughley, R.J. 1987. Acacias and their root nodule bacteria. *In J.W. Turnbull, ed., Australian acacias in developing countries*, pp. 45–49. ACIAR, Canberra, Australia.
139. Rowe, E.C., Hairiah, K., Giller, K.E., Noordwijk, M. van, and Cadisch, G. 1999. Testing the safety-net role of hedgerow tree roots by ¹⁵N placement at different soil depths. *Agroforestry Systems* 43:81–93.
140. Samba, R.T., Lajudie, P. de, Gillis, M., Neyra, M., Barreto, M.M.S., and Dreyfus, B. 1999. Diversity of rhizobia nodulating *Crotalaria* spp. from Senegal. *Symbiosis* 27:259–268.
141. Sanginga, N. 1988. Nodulation and growth of some nitrogen fixing trees in relation to nutrient levels and *Rhizobium* in Nigeria, Zaire and Zimbabwe. *Nitrogen Fixing Tree Research Reports* 6: 3–5.
142. Saur, E., S. Carcella, S. Guezennec, and A. Rousteau. 2000. Nodulation of legume species in wetlands of Guadeloupe (Lesser Antilles). *Wetlands* 20(4): 730–734.
143. Selvaraj, K., and Abragam, D.A. 1999. Occurrence of Ca-oxalate crystals in nitrogen-fixing root nodules and other tissues of some tropical legumes. *Phytomorphology* 49(1): 79–87.
144. Setiadi, D., McKinnell, F.H., Harwood, C.E., and Dart, P. 1997. Survival and growth of Australian *Acacia* species and their response to rhizobium inoculation in field trials on two contrasting soil types in Indonesia. *Forest, Farm, and Community Tree Research Reports* 2:29–34.
145. Shaw, J.E., Reynolds, T., and Sprent, J.I. 1997. A study of the symbiotic importance and location of *nod* gene inducing compounds in two nodulating and two non-nodulating tropical tree species. *Plant and Soil* 188:77–82.
146. Smith, R.S., S. Thome, and R. Randall. 1988. *Rhizobia* inoculants for various leguminous species. *The Nitragin Company Technical Bulletin* 102: 1–8.
147. Sparrow, S.D., Jr. 1986. Effects of inoculation on nodulation, growth and nitrogen accumulation by seedling-year alfalfa, *Medicago media* Persoon, in three subarctic soils. *Crop Res. (Hort. Res.)* 25: 133–142.
148. Sprent, J.I. 1988. Problems and potentials for nitrogen fixation in deserts. *In E.E. Whitehead, C.F. Hutchinson, and B.N. Timmerman, eds., Arid lands: today and tomorrow*, pp. 1049–1062. Westview Press Inc., Coulter, Colorado.
149. Sprent, J.I. 2000. Nodulation as a taxonomic tool. *In P.S. Herendeen and A. Bruneau, eds., Advances in Legume Systematics* 9, pp. 21–44. Royal Botanic Gardens, Kew, England.
150. Sprent, J.I. 2001. Nodulation in legumes. Royal Botanic Gardens, Kew.
151. Sprent, J.I. 1995. Legume trees and shrubs in the tropics: N₂ fixation in perspective. *Soil Biology and Biochemistry* 27(4/5):401–407.

152. Sprent, J.I., Geoghegan, I.E., and Whitty, P.W. 1996. Natural abundance of ^{15}N and ^{13}C in nodulated legumes and plants in the cerrado and neighbouring regions of Brazil. *Oecologia* 105:440-446.
153. Strijdom, B.W. 1998. South African studies on biological nitrogen-fixing systems and the exploitation of the nodule bacterium-legume symbiosis. *South African Journal of Science* 94:11-23.
154. Sylvester-Bradley, R., L.A. de Oliveira, J.A. de Podesta Filho, and T.V. St. John. 1980. Nodulation of legumes, nitrogenase activity of roots and occurrence of nitrogen-fixing *Azospirillum* spp. In representative soils of central Amazonia. *Agro-Ecosystems* 6: 249-267.
155. Tan, Z.-Y., Wang, E.-T., Peng, G.-X., Zhu, M.-E., Martínez-Romero, E., and Chen, W.-X. 1999. Characterization of bacteria isolated from wild legumes in the north-western regions of China. *International Journal of Systematic Bacteriology* 49:1457-1469.
156. Tan, Z.Y., Zhu, M.E., Wang, E.T., Cheng, L.J., Li, Y., Ye, Q., Li, Z.F., and Chen, W.X. 1996. Characterization of rhizobial strains isolated from the loess plateau of China. In F.D. Li, T.A. Lie, W.X. Chen, and J.C. Zhou, eds., *Diversity and Taxonomy of Rhizobia*, pp. 85-97. China Agricultural Sciencetech Press, Beijing, China.
157. Thomas, B.D., and Bowman, W.D. 1998. Influence of N_2 -fixing *Trifolium* on plant species composition and biomass production in alpine tundra. *Oecologia* 115:26-31.
158. Tisserant, B., Brenac, V., Requena, N., Jeffries, P., and Dodd, J.C. 1998. The detection of *Glomus* spp. (arbuscular mycorrhizal fungi) forming mycorrhizas in three plants, at different stages of seedling development, using mycorrhiza-specific isozymes. *New Phytologist* 138:225-239.
159. Turvey, N.D., and P.J. Smethurst. 1983. Nitrogen fixing plants in forest plantation management. In J.C. Gordon and C.T. Wheeler, eds., *Biological nitrogen fixation in forest ecosystems: foundations and applications*, pp. 233-257. Martinus Nijhoff/Dr. W. Junk Publishers, The Hague.
160. Vielma A., M. 1999. Caracterización de cepas autóctonas de *Bradyrhizobium* sp. aisladas de *Lupinus* spp. *Revista de la Facultad de Agronomía, Universidad del Zulia* 16:495-508.
161. Wang, E.-T., Rogel, M.A., García-de los Santos, A., Martínez-Romero, J., Cevallos, M.A., and Martínez-Romero, E. 1999. *Rhizobium etli* bv. mimosae, a novel biovar isolated from *Mimosa affinis*. *International Journal of Systematic Biology* 49:1479-1491.
162. Wang, S.Y., and Chen, W.X. 1996. Numerical taxonomy and DNA relatedness of rhizobia isolated from *Astragalus* spp. In F.D. Li, T.A. Lie, W.X. Chen, and J.C. Zhou, eds., *Diversity and Taxonomy of Rhizobia*, pp. 79-84. China Agricultural Sciencetech Press, Beijing, China.
163. Wester, J., and P. Höberg. 1989. New nodulating legume tree species from Guinea-Bissau, West Africa. *Forest Ecology and Management* 29: 311-314.
164. Wijesundra, T.I.L., Van Holm, L.H.J., and Kulasoorya, S.A. 2000. Rhizobiology and nitrogen fixation of some tree legumes native to Sri Lanka. *Biology and Fertility of Soils* 30:535-543.
165. Workman, S.W. 1986. Nodulation trials with selected arboreal legumes of Burkina Faso, West Africa. *Nitrogen Fixing Tree Research Reports* 4: 7-8.
166. Wynne, J.C., G.H. Elkan, T.J. Schneeweis, T.G. Isleib, C.M. Preston, and C.A. Meisner. 1978. Increasing nitrogen fixation of the peanut. *Proc. Amer. Peanut Res. Educ. Assoc.* 10(1): 22-29.
167. Yates, R.J., Howieson, J.G., O'Hara, G.W., and Gintzburger, G. 2000. The interaction of non-indigenous perennial legumes from Mediterranean-type climates with root-nodule bacteria in soils of the north-west of Western Australia. In D. Eldridge and D. Freudenberger, eds., *People and rangelands: building the future*, Proceedings of the VI International Rangeland Congress, Townsville, Queensland, Australia, 19-23 July 1999, pp. 321-322. VI International Rangeland Congress, Aitenvale, Queensland.
168. Yoneyama, T. 1987. N_2 fixation and natural ^{15}N abundance of leguminous plants and *Azolla*. *Bull. Natl. Inst. Agrobiol. Resour.* 3: 59-87.
169. Zeng, Y., Lui, S.G., Holger, B., and Bai, L.H. 1996. The temperature, pH, ultraviolet and salt tolerance of rhizobia strains from grasslands on Qinghai-Tibet plateau. In F.D. Li, T.A. Lie, W.X. Chen, and J.C. Zhou, eds., *Diversity and Taxonomy of Rhizobia*, pp. 301-307. China Agricultural Sciencetech Press, Beijing, China.
170. Zhang, F.-Q., Chen, Z.-X., Searle, S.D., Li, X.-M., Zhou, J.-J., and Li.Q. 1998. Temperate Australian *Acacia* species elimination trials in southern China. In J.W. Turnbull, H.R. Crompton, and K. Pinyopusarerk, eds., *Recent Developments in Acacia Planting: Proceedings of an international workshop held in Hanoi, Vietnam, 27-30 October 1997*, pp. 36-44. Australian Centre for International Agricultural Research, Canberra, Australia.

Leguminous Genera of Unknown Response to Rhizobia

Harold D.L. Corby and Joe.H. Kirkbride, Jr.

These lists of leguminous genera whose response to rhizobia is still unknown are presented to encourage their exploration. They are arranged geographically in the following six tables so that *Bean Bag* Readers may see at a glance the task awaiting them on their doorsteps.

Response has been reported for 493 legume genera comprising more than 17,450 species, leaving 195 genera, comprising some 549 species, still to be explored. One hundred and two genera, 52.6%, are monotypic; 86, 43.9%, have 2–10 species; and six, 3.1%, have 11–20 species. The vast majority is found in tropical areas, and only two genera, *Adenopodia* C. Presl and *Pomaria* Cav., occur in both the New and Old Worlds. *Pomaria* is also the largest genus with approximately 20 species. The genera, as understood here, are those accepted by the US Department of Agriculture in the Germplasm Resources Information Network [GRIN] (USDA 2001). The tribal classification is that of Polhill (1994a, 1994b), updated to reflect subsequent work (Kirkbride *et al.* 2000; USDA loc. cit.). The number of species per genus are those of Polhill and Raven (1981), updated to accord with those of Kirkbride *et al.* (loc. cit.).

Genera are listed by the mid-latitude of their distribution as given in Polhill and Raven (loc. cit.), updated to accord with Kirkbride *et al.* (loc. cit.). Genera occurring continuously across the equator are listed as *equatorial*, regardless of their extent on either side of it; those occurring on both sides of the equator, but not across it, are listed by the mid-latitudes of their northern and southern distributions.

Nodule Hunting

For those who may be unfamiliar with nodule hunting, it may be said that by far the best method of search is to raise seedlings in coarse sharp sand, with an admixture of soil from the rhizosphere of flourishing plants of the species concerned. Then, when seedlings are at least two months old, or at flowering if that comes sooner, their roots should be washed clean of soil to reveal any nodules present.

Those seeking nodules in the field must, when exploring root-systems too big to be dug out in their entirety, always trace the nodule-bearing roots back to the plant itself—only thus may the nodule be surely related to its supposed parent.

Effective rhizobial nodules (those fixing nitrogen—those being sought) may usually be recognized in being lateral outgrowths of the root with reddish-pink cores. However, these two criteria are not infallible, so specimens of any nodule found should be dried over silica gel for later formal identification; at the same time, specimens of the parent plant should be pressed and dried for the same purpose.

With seemingly non-nodulating species, it is suggested that successive batches of seedlings be raised, using a bacterial inoculum from a different source with each batch, until the tally of flourishing plants without nodules reaches at least 100. And then, and only then, to report that the species has been found to be non-nodulating! Fuller treatment of this subject may be found in Corby (1981, 1988) and Sprent (2001).

Table 1. African genera whose response to rhizobia is unknown and their distribution in Africa.

Genus	Tribe	No. of Species	Distribution
30° or more North			
<i>Argyrocytisus</i> (Maire) Frodin & Heywood ex Raynaud	Genisteae	1	Atlas Mountains
<i>Dorycnopsis</i> Boiss.	Loterae	1	E Mediterranean, N Morocco
<i>Erinacea</i> Adans.	Genisteae	1	NW Africa (and SW Europe)
<i>Spartidium</i> Pomel	Crotalarieae	1	North Africa
10–29° North			
<i>Clitoriopsis</i> R. Wilczek	Phaseoleae	1	Zaire, Sudan
<i>Distemonanthus</i> Benth.	Cassieae	1	West Tropical Africa
<i>Griffonia</i> Baill.	Cercideae	4	West Tropical Africa
<i>Melliniella</i> Harms	Desmodieae	1	West Tropical Africa
10–29° North contd.			
<i>Platycelyphium</i> Harms	Sophoreae	1	E and NE Africa
<i>Talbotiella</i> Baker f.	Detarieae	3	West Africa
<i>Vernifrux</i> J. B. Gillett	Loteae	1	N Ethiopia, Sudan, Eritrea, Yemen

3–9° North			
<i>Airyantha</i> Brummitt	Sophoreae	2	Guinea-Congo forests (and Borneo 1)
<i>Aphanocalyx</i> Oliv.	Detarieae	14	Guinea-Congo forests and Sierra Leone—Liberia—Ivory Coast
<i>Aubrevillea</i> Pellegr.	Mimoseae	2	Guinea-Congo forests
<i>Chidlowia</i> Hoyle	Caesalpinieae	1	Sierra Leone to Ghana
<i>Dicraeopetalum</i> Harms	Sophoreae	3	Kenya, Ethiopia (and Madagascar)
<i>Gilletiodendron</i> Vermeesen	Detarieae	5	Guinea-Congo forests
<i>Haplormosia</i> Harms	Sophoreae	1	Sierra Leone—Gabon
<i>Lebruniodendron</i> J. Léonard	Detarieae	1	Forests of Gulf of Guinea
<i>Leonardoxa</i> Aubrev.	Detarieae	1	Guinea-Congo forests
<i>Oddoniodendron</i> De Wild.	Detarieae	2	Gulf of Guinea forests
<i>Oxystigma</i> Harms	Detarieae	5	Guinea-Congo forests
<i>Pellegriniodendron</i> J. Léonard	Detarieae	1	Guinea-Congo forests
<i>Plagiosiphon</i> Harms	Detarieae	5	Guinea-Congo forests
<i>Polystemonanthus</i> Harms	Detarieae	1	Liberia and Ivory Coast
<i>Stemonocoleus</i> Harms	Detarieae	1	Guinea-Congo forests
<i>Tessmannia</i> Harms	Detarieae	11	Guinea-Congo and East Africa
Equatorial Africa			
<i>Adenopodia</i> C. Presl	Mimoseae	10	Africa 4 (and Central America and Mexico 3, Amazonia 3)
<i>Bikinia</i> Wieringa	Detarieae	10	W central Africa
<i>Bussea</i> Harms	Caesalpinieae	6	Tropical Africa (and Madagascar)
<i>Camoensia</i> Welw. ex Benth. & Hook. f.	Sophoreae	2	Forests of Gulf of Guinea
<i>Eurypetalum</i> Harms	Detarieae	3	Cameroon—Gabon
<i>Fillaeopsis</i> Harms	Mimoseae	1	Nigeria—Zaire—Angola
<i>Nesphostylis</i> Verdc.	Phaseoleae	3	Africa 1 (and India 1, Burma 1)
<i>Normandiodendron</i> J. Léonard	Detarieae	2	Zaire, Gabon, Cabinda
<i>Ostryocarpus</i> Hook. f.	Millettieae	6	Tropical Africa to SE Asia
<i>Pachyelasma</i> Harms	Caesalpinieae	1	Nigeria, Zaire, Gabon, Central African Republic
<i>Paracalyx</i> Ali	Phaseoleae	6	Africa to Asia
<i>Physostigma</i> Balf.	Phaseoleae	4	Tropical Africa
<i>Platysepalum</i> Welw. ex Baker	Millettieae	12	Tropical Africa
<i>Pseudoeriosema</i> Hauman	Phaseoleae	6	Tropical Africa
<i>Pseudoprosopis</i> Harms	Mimoseae	4	Tropical Africa
<i>Schefflerodendron</i> Harms	Millettieae	6	Tropical Africa
<i>Spathionema</i> Taub.	Phaseoleae	1	Tropical E Africa
<i>Stachyothyrsus</i> Harms	Caesalpinieae	3	Sierra Leone—Zaire—Gabon
<i>Vatovaea</i> Chiov.	Phaseoleae	1	Africa
3–9° South			
<i>Augouardia</i> Pellegr.	Detarieae	1	Gabon
<i>Englerodendron</i> Harms	Detarieae	1	Usambara Mountains, Tanzania
<i>Librevillea</i> Hoyle	Detarieae	1	Gabon
<i>Michelsonia</i> Hauman	Detarieae	1	Zaire forests
<i>Neochevalierodendron</i> J. Léonard	Detarieae	1	Gabon
<i>Pseudomacrolobium</i> Hauman	Detarieae	1	Forests of Congo basin
<i>Sindoropsis</i> J. Léonard	Detarieae	1	Gabon
<i>Stuhlmannia</i> Taub.	Caesalpinieae	1	Coastal forests of Tanzania

10–21° South			
<i>Adenolobus</i> (Harv. ex Benth. & Hook. f.) Torre & Hillc.	Cercideae	2	SW Africa
<i>Carrissoa</i> Baker f.	Phaseoleae	1	Angola
<i>Icuria</i> Wieringa	Detarieae	1	Coastal dunes of Mozambique
<i>Rhynchotropis</i> Harms	Indigofereae	2	South central Africa
22° or more South			
<i>Chrysoscias</i> E. Mey.	Phaseoleae	6	South Africa
<i>Polhillia</i> C.H. Stirt.	Genisteae	5	Cape Province, South Africa
<i>Pomaria</i> Cav.	Caesalpinieae	20	South Africa (and SW USA, N Mexico, South America)
<i>Stirtonanthus</i> B.-E. van Wyk & Schutte	Podalyriinae	3	Cape Province, South Africa
<i>Umtiza</i> Sim	Detarieae	1	South Africa

Ed. Note. Molecular data places *Umtiza* within subfamily Caesalpinioideae s.l but excludes it from Detarieae s.l.

Table 2. Madagascan genera whose response to rhizobia is unknown.

Genus	Tribe	No. of Species	Distribution
<i>Alantsilodendron</i> Villiers	Mimoseae	8	
<i>Baudouinia</i> Baill.	Cassieae	4	
<i>Brandzeia</i> Baill.	Detarieae	1	
<i>Brenierea</i> Humbert	Cercideae	1	
<i>Dicraeopetalum</i> Harms	Sophoreae	3	Ethiopia to N Kenya
<i>Disynstemon</i> R. Vig.	Millettieae	1	
<i>Eligmocarpus</i> Capuron	Cassieae	1	
<i>Gagnebina</i> Neck. ex DC.	Mimoseae	2	Comoros, Mauritius, Reunion
<i>Lemurodendron</i> Villiers & P. Guinet	Mimoseae	1	
<i>Lemuropisum</i> H. Perrier	Caesalpinieae	1	
<i>Mendoravia</i> Capuron	Cassieae	1	
<i>Neoapaloxylon</i> Rauschert	Detarieae	2	
<i>Neoharmsia</i> R. Vig.	Sophoreae	2	
<i>Ormocarpopsis</i> R. Vig.	Aeschynomeneae	5	
<i>Peltiera</i> Du Puy & Labat	Aeschynomeneae	2	
<i>Phylloxylon</i> Baill.	Indigofereae	5	
<i>Pongamiopsis</i> R. Vig.	Millettieae	2	
<i>Pyranthus</i> Du Puy & Labat	Millettieae	6	
<i>Sakoanala</i> R. Vig.	Sophoreae	2–3	
<i>Sylvichadsia</i> Du Puy & Labat	Millettieae	4	
<i>Tetrapterocarpon</i> Humbert	Caesalpinieae	1	
<i>Vaughania</i> S. Moore	Indigofereae	11	

Table 3. New World genera whose response to rhizobia is unknown and their distribution in the Americas and Caribbean islands.

Genus	Tribe	No. of Species	Distribution
30° or more North			
<i>Errazurizia</i> Phil.	Amorpheae	4	Arizona, Baja California, Sonora
<i>Kanaloa</i> Lorence & K. R. Wood	Mimoseae	1	Kaho’olawe, Hawaii
<i>Lackeya</i> Fortunato et al.	Phaseoleae	1	SE USA
<i>Peteria</i> A. Gray	Robinieae	4	Mexico, USA

30° or more North contd.			
<i>Pomaria</i> Cav.	Caesalpinieae	20	SW USA, N Mexico (and South America, South Africa)
<i>Rupertia</i> J. W. Grimes	Psoraleeae	3	SW Canada, W USA and N Baja California, Mexico
<i>Sphinctospermum</i> Rose	Robinieae	1	Mexico, USA
20–29° North			
<i>Behaimia</i> Griseb.	Millettieae	1	Cuba
<i>Calliandropsis</i> H. M. Hern. & P. Guinet	Mimoseae	1	Central Mexico
<i>Genistidium</i> I. M. Johnst.	Robinieae	1	Mexico and Texas, USA
<i>Herpyza</i> C. Wright	Phaseoleae	1	Cuba
<i>Hesperothamnus</i> Brandegees	Millettieae	5	Mexico
<i>Painteria</i> Britton & Rose	Ingeae	3	Mexican plateau
<i>Pictetia</i> DC.	Aeschynomeneae	5	Cuba
<i>Pomaria</i> Cav.	Caesalpinieae	20	SW USA, N Mexico (and South America, South Africa)
<i>Prosopidastrum</i> Burkart	Mimoseae	1	Mexico (and Patagonia 1)
10–19° North			
<i>Adenopodia</i> C. Presl	Mimoseae	10	Mexico and Central America 3, Amazonia 3 (and Africa 4)
<i>Apoplanesia</i> C. Presl	Amorpheae	1	Mexico—Venezuela
<i>Arcoa</i> Urb.	Caesalpinieae	1	Santo Domingo
<i>Guinetia</i> L. Rico & M. Sousa	Ingeae	1	Oaxaca, Mexico
<i>Hybosema</i> Harms	Robinieae	2	S Mexico, N Central America
<i>Lennea</i> Klotzsch	Robinieae	5	Central America, Mexico
<i>Myrospermum</i> Jacq.	Sophoreae	2	S Texas, USA to northern South America
<i>Pictetia</i> DC.	Aeschynomeneae	6	West Indies
<i>Pomaria</i> Cav.	Caesalpinieae	20	SW USA, N Mexico (and South America, South Africa)
<i>Prioria</i> Griseb.	Detarieae	1	Panama, Costa Rica, Jamaica, Colombia, Central America
<i>Rhodopis</i> Urb.	Phaseoleae	1	West Indies
<i>Sphinga</i> Barneby & J. W. Grimes	Ingeae	3	S Mexico, N. Guatemala, Cuba, N coastal Colombia and Venezuela
<i>Uribea</i> Dugand & Romero	Sophoreae	1	Costa Rico, Colombia
3–9° North			
<i>Brachycylix</i> (Harms) R. S. Cowan	Detarieae	1	Colombia
<i>Candolleodendron</i> R. S. Cowan	Swartzieae	1	French Guiana, mouth of Amazon
<i>Fissicalyx</i> Benth.	Dalbergieae	1	Venezuela, Guyana, Panama
<i>Margaritolobium</i> Harms	Millettieae	1	Margarita Island, Venezuela
<i>Orphanodendron</i> Barneby & J. W. Grimes	Caesalpinieae	1	Antioquia, Colombia
<i>Spirotropis</i> Tul.	Sophoreae	1	NE South America
Equatorial America			
<i>Adenopodia</i> C. Presl	Mimoseae	10	Mexico and Central America 3, Amazonia 3 (and Africa 4)
<i>Apurimacia</i> Harms	Millettieae	4	Tropical and subtropical South America
<i>Browneopsis</i> Huber	Detarieae	6	Panama, Colombia, Ecuador, Peru
<i>Collaea</i> DC.	Phaseoleae	3	Tropical and subtropical South America
<i>Cymbosema</i> Benth.	Phaseoleae	1	Mexico—Amazonia
<i>Ecuadendron</i> D. A. Neill	Detarieae	1	W Ecuador
<i>Jacqueshuberia</i> Ducke	Caesalpinieae	3	Brazil, Colombia, Venezuela, Peru

Equatorial America contd.			
<i>Muelleria</i> L. f.	Millettieae	2	Tropical America
<i>Paloue</i> Aubl.	Detarieae	4	Tropical America
<i>Paloveopsis</i> R. S. Cowan	Detarieae	1	NE South America
<i>Soemmeringia</i> Mart.	Aeschynomeneae	1	Brazil, Venezuela, Bolivia
3–9° South			
<i>Androcalymma</i> Dwyer	Cassieae	1	Amazonia
<i>Uleanthus</i> Harms	Sophoreae	1	Amazon basin
<i>Weberbauerella</i> Ulbr.	Aeschynomeneae	2	Peru
10–29° South			
<i>Blanchetiodendron</i> Barneby & J. W. Grimes	Ingeae	1	NE Brazil
<i>Cascaronia</i> Griseb.	Dalbergieae	1	Bolivia, Argentina
<i>Fiebrigiella</i> Harms	Aeschynomeneae	1	Bolivia, Peru
<i>Leucochloron</i> Barneby & J. W. Grimes	Ingeae	4	E Brazil
<i>Mysanthus</i> G. P. Lewis & A. Delgado	Phaseoleae	1	Bahia, Brazil
<i>Oryxis</i> A. Delgado & G. P. Lewis	Phaseoleae	1	Minas Gerais, Brazil
<i>Panurea</i> Spruce ex Benth. & Hook. f.	Sophoreae	1	Colombia, Brazil
<i>Petaladenium</i> Ducke	Sophoreae	1	Brazil
<i>Piptadeniopsis</i> Burkart	Mimoseae	1	Paraguay
<i>Pomaria</i> Cav. <i>Pomaria</i> Cav. contd.	Caesalpinieae	20	South America (and SW USA, N Mexico, South Africa)
<i>Sellocharis</i> Taub.	Genisteae	1	SE Brazil
<i>(Thylacanthus</i> Tul.	Detarieae	1	Brazil) see Ed. Note.
30° or more South			
<i>Anarthrophyllum</i> Benth.	Genisteae	15	Chile, Argentina
<i>Pomaria</i> Cav.	Caesalpinieae	20	South America (and SW USA, N Mexico, South Africa)
<i>Prosopidastrum</i> Burkart	Mimoseae	1	Patagonia (and Mexico 1)
<i>Ramorinoa</i> Speg.	Dalbergieae	1	Argentina
<i>Stenodrepanum</i> Harms	Caesalpinieae	1	Argentina, Brazil
<i>Zuccagnia</i> Cav.	Caesalpinieae	1	Argentina, Chile

Ed .Note. *Thylacanthus* Tul. , thought to be a monotypic genus from Brazil based on a single collection is *Julbernardia paniculata* (Benth.) Troupin from Mozambique (Breteler in comm. G.P.Lewis, 1995).

Table 4. Asian and Indo-Pacific island genera whose response to rhizobia is unknown and their distribution in the Asia and the Indo-Pacific islands.

Genus	Tribe	No. of Species	Distribution
40° or more North			
<i>Ammodendron</i> Fisch. ex DC.	Sophoreae	6	Central Asia
<i>Eversmannia</i> Bunge	Hedysareae	1	Caspian Sea to Tien Shan
<i>Paracalyx</i> Ali	Phaseoleae	6	Asia (and Africa)
30–39° North			
<i>Craspedolobium</i> Harms	Millettieae	1	W China
<i>Diphyllarium</i> Gagnep.	Phaseoleae	1	Indochina, Indonesia, Philippines
<i>Oreophysa</i> (Bunge ex Boiss.) Bomm.	Galegeae	1	Mountains of N Iran
<i>Salweenia</i> Baker f.	Sophoreae	1	Tibet
<i>Sarcodum</i> Lour.	Millettieae	2	SE Asia, Japan (Solomon Is. 1, Phillipines,1)

30–39° North contd.			
<i>Sinodolichos</i> Verdc.	Phaseoleae	2	China and Burma
<i>Spongiocarpella</i> Yakovlev & N. Ulziykh.	Galegeae	9	Central China, foothills of the Himalayas, Mongolia
<i>Stracheya</i> Benth.	Hedysareae	1	N India, Nepal, Bhutan, SW China
3–29° North			
<i>Afgekia</i> Craib	Millettieae	3	S China, Burma, Thailand
<i>Airyantha</i> Brummitt	Sophoreae	1	Borneo (and Africa)
<i>Antheroporum</i> Gagnep.	Millettieae	2	Thailand, Indochina, S China
<i>Burkilliodendron</i> Sastry	Millettieae	1	Perak, Malaysia
<i>Cochlianthus</i> Benth.	Phaseoleae	2	Himalayas, SW China
<i>Cruddasia</i> Prain	Phaseoleae	1	Burma, Thailand, S China
<i>Desmodiastrum</i> (Prain) A. Pramanik & Thoth.	Desmodieae	2	S India, Indochina, E Java
<i>Eleiotis</i> DC.	Desmodieae	2	India
<i>Endertia</i> Steenis & de Wit	Detarieae	1	Borneo
<i>Endosamara</i> R. Geesink	Millettieae	1–2	S India to Philippines
<i>Luzonia</i> Elmer	Phaseoleae	1	Philippines, Indonesia
<i>Nesphostylis</i> Verdc.	Phaseoleae	3	India 1, Burma 1 (and Africa 1)
<i>Nogra</i> Merr.	Phaseoleae	3	India, Thailand, China
<i>Ostryocarpus</i> Hook. f.	Millettieae	6	Tropical Africa to SE Asia
<i>Pseudosindora</i> Symington	Detarieae	1	Borneo, Sarawak
<i>Spongiocarpella</i> Yakovlev & N. Ulziykh.	Galegeae	9	Central China, foothills of the Himalayas, Mongolia
<i>Storckiella</i> Seem.	Cassieae	3	New Caledonia and Fiji
<i>Stracheya</i> Benth.	Hedysareae	1	N India, Nepal, Bhutan, SW China
Equatorial Asia			
<i>Fordia</i> Hemsl.	Millettieae	10	Continental SE Asia, Sumatra, Borneo
<i>Kingiodendron</i> Harms	Detarieae	6	India, East Indies
<i>Kunstleria</i> Prain	Millettieae	7	Malaya, Borneo, Sumatra, Philippines, New Guinea, S India
<i>Leucostegane</i> Prain	Detarieae	2	Malaya, Sarawak
<i>Macropsychanthus</i> Harms ex K. Schum. & Lauterb.	Phaseoleae	4	Philippines, New Guinea, Micronesia
<i>Mecopus</i> Benn.	Desmodieae	1	India to S China and Malayan Islands
<i>Neocollettia</i> Hemsl.	Desmodieae	1	Burma, Java
<i>Serianthes</i> Benth.	Ingeae	10	Malay peninsula and archipelago
<i>Sympetalandra</i> Stapf	Caesalpinieae	5	East Indies
<i>Trifidacanthus</i> Merr.	Desmodieae	2	Vietnam, Hainan, Philippines, Lesser Sunda Islands
3° or more South			
<i>Arthroclianthus</i> Baill.	Desmodieae	10	New Caledonia
<i>Desmodiastrum</i> (Prain) A. Pramanik & Thoth.	Desmodieae	2	S India, E Java
<i>Kalappia</i> Kosterm.	Cassieae	1	Celebes
<i>Nephrodesmus</i> Schindl.	Desmodieae	6	New Caledonia
<i>Sarcodum</i> Lour.	Millettieae	2	Solomon Islands 1, Philippines 1 (and SE Asia and Japan)
<i>Storckiella</i> Seem.	Cassieae	3	New Caledonia, Fiji

Table 5. Australian genera whose response to rhizobia is unknown and their distribution in Australia.

Genus	Tribe	No. of Species	Distribution
<i>Aenictophyton</i> A. T. Lee	Bossiaeeae	1	Northern Terr. and Western Australia
<i>Austrodolichos</i> Verdc.	Phaseoleae	1	Australia
<i>Austroteenisia</i> R. Geesink	Millettieae	4	E coastal Australia
<i>Jansonia</i> Kippist	Mirbelieae	1	SW Western Australia
<i>Labichea</i> Gaudich. ex DC.	Cassieae	9	Australia
<i>Pararchidendron</i> I. C. Nielsen	Ingeae	2	Australia (and Asia)
<i>Phylacium</i> Benn.	Desmodieae	2	N Queensland (and Asia)
<i>Plagiocarpus</i> Benth.	Bossiaeeae	1	N Australia
<i>Ptychosema</i> Benth. ex Lindl.	Bossiaeeae	2	Central and Western Australia
<i>Streblorrhiza</i> Endl.	Carmichaelieae	1	Philip Island
<i>Urodon</i> Turcz.	Mirbelieae	1–3	SW Australia

Table 6. European and Mediterranean genera whose response to rhizobia is unknown and their distribution in Europe and Mediterranean basin.

Genus	Tribe	No. of Species	Distribution
<i>Cytisopsis</i> Jaub. & Spach	Loteae	2	E Mediterranean and North Africa
<i>Dorycnopsis</i> Boiss.	Loteae	1	E Mediterranean, N Morocco
<i>Echinospartum</i> (Spach) Fourr.	Genisteae	3	SW Europe
<i>Erinacea</i> Adans.	Genisteae	1	SW Europe and NW Africa
<i>Gonocytisus</i> Spach	Genisteae	3	E Mediterranean
<i>Hammatolobium</i> Fenzl	Loteae	2	Mediterranean basin
<i>Podocytisus</i> Boiss. & Heldr.	Genisteae	1	Balkan peninsula, W Turkey
<i>Sartoria</i> Boiss. & Heldr.	Hedysareae	1	S Turkey

References

Corby, H.D.L. 1981. The systematic value of leguminous root nodules. In R.M. Polhill and P.H. Raven, eds. *Advances in Legume Systematics*, Part 2, pp. 657–669. London, Royal Botanic Gardens, Kew.

Corby, H.D.L. 1988. Types of Rhizobial nodules and their distribution among the Leguminosae. *Kirkia* 13: 53–123.

Kirkbride, J.H., Jr., and Gilbert, L.A. 2001. *Legume (Fabaceae) nodulation*. USDA Technical Bulletin, in press.

Kirkbride, J.H., Jr., Gunn, C.R., Weitzman, A.L., and Dallwitz, M.J. 2000. *Legume (Fabaceae) Fruits and Seeds*, CDROM. Boone, North Carolina, Parkway Publishers.

Polhill, R.M. 1994a. Classification of the Leguminosae. In F.A. Bisby, J. Buckingham, and J.B. Harborne, eds., *Phytochemical Dictionary of the Leguminosae*, pp. xxxv–xlvi. Chapman & Hall, London, Enngland.

Polhill, R.M. 1994b. Complete synopsis of legume genera. In F.A. Bisby, J. Buckingham, and J.B. Harborne, eds., *Phytochemical Dictionary of the Leguminosae*, pp. xlix–lvii. Chapman & Hall, London, Enngland

Polhill, R.M., and Raven, P.H., eds. 1981. *Advances in Legume Systematics*, Part 1. London, Royal Botanic Gardens, Kew.

Sprent, J.I. 2001. *Nodulation in Legumes*. London, Royal Botanic Gardens, Kew.

USDA, ARS, National Genetic Resources Program. 2001. Germplasm Resources Information Network - (GRIN), GRIN Taxonomy. [Online Database] National Germplasm Resources Laboratory, Beltsville, Maryland. Available: <http://www.ars-grin.gov/npgs/tax> (30 August 2001)

GLEANINGS

Ed. Note: Names in all capital letters are *Bean Bag* Readers. Their full names and addresses are listed in the *Bean Bag* Current Readers' Directory soon to be available on the www server of the Royal Botanic Gardens, Kew, United Kingdom. See "From the Editor" for the URL.

ALBUQUERQUE is studying medicinal legumes, *Zornia reticulata* Smith, *Mimosa pudica* L. and *Copaifera* spp. and would like any information about these (and other) tropical legumes used in popular medicine. Offers information about tropical legumes of Amazonia – José Maria de Albuquerque, Caixa Postal 716, 66.017-970-Belém-Pará, Brasil.

ARAMBARRI A.M., Stenglein, S.A., Colares, M.N., Novoa, M.C. and Vizcaino, C.E. are working on "Leaf epidermal characteristics in Old and New World species of *Lotus* (Fabaceae: Loteae)". Their purpose is to discover diagnostic characters for future studies.

BEYRA-MATOS continues working on tribe Phaseoleae and defended her PhD thesis on *Pictetia* (PhD thesis) in November 2000 at Havana University, which was approved by the tribunal of scientific grade at Havana University.

CACCAVARI is undertaking a pollen study of American genera of the Piptadenia Group: *Adenopodia*, *Stryphnodendron*, *Mimosa*, *Piptadeniopsis*, *Microlobius*, *Piptadenia*, *Pseudopiptadenia* and *Parapiptadenia*. Flora palinologica del Nordeste Argentino. Needs pollen material of species of the different American taxa of the Piptadenia Group and offers pollen slides of Mimosoideae of Argentina.

FORERO has been up-dating the Mimosoideae in COL (National Herbarium of Colombia) based on monographs by Barneby & Grimes and by Pennington. Studies on the Mimosoideae of Colombia are underway.

HIRSCH is working on genetics of sweet clover (*Melilotus albus* Medikus) and needs Rhizobium inocula for *Gleditsia*. She offers *M. albus* mutants.

KIAM is working on nodulation of legumes in Cameroonian forest zone. Needs lab material for the study of root nodule or any information on nodulation of legumes and offers seed of some Cameroonian legumes such as *Pterocarpus* and *Erythrophleum*.

PAVLOVA needs legume research material of *Bauhinia* L.: *B. aculeata* L., *B. acuminata* L., *B. cumanensis* H.B.K., *B. diphylla* Ham., *B. galpinii* Br., *B. hookeri* Muell., *B. jumanensis* Fr., *B. monandra* Kurz., *B. purpurea* L., *B. racemosa* Lam., *B. rufescens* Lam. and especially material of *B. tomentosa* L., *B. variegata* L. For each species 20–30 seeds and/or a few coloured photographs of the plants in flower are needed.

PIERGIOVANNI with Nenno is maintaining a germplasm database on Italian landraces of *Phaseolus vulgaris* L which is available on <http://www.ba.cnr.it/~germap14/ilcb/>. They offer and need seed material of the same.

PISTRICK needs land races and old varieties of cultivated legumes and its wild relatives. Offers legumes from the Gatersleben genebank (cp. Index seminum). <http://fax-serve.ipk-gatersleben.de>

PLANCHUELO is Director of a new Economic Botany project that combines studies in seed anatomy and morphology of crop legumes with taxonomic studies of wild species. Needs Herbarium specimens of *Lupinus* and *Crotalaria* and offers Herbarium specimens of legumes from Central Argentina.

ROBILLARD offers legume publications The list of titles is available from Library-res@em.agr.ca.

SARAVANAKUMAR is working on comparative studies of proteins, chromosome and DNA of some populations of *Vigna* spp. Needs material of grain legumes and its wild relatives and information concerning genetics and conservation of grain legumes and its wild relatives. Offering wild relatives of grain legumes literature/seeds such as *Vigna* and *Cajanus*.

TUCKER is working on comparative floral ontogeny of Swartzieae and needs young floral inflorescences and buds of Swartzia, other taxa in Swartzieae. Offers reprints of publications on floral development in various legumes.

VANDERBORGHT is maintaining a Phaseoleae – Phaseolinae collections chiefly centred on wild *Phaseolus* and *Vigna* species. Offers and needs seed material of the same. List of the taxa included in the collection is available on <http://www.bn.fgov.be/research/collections/living/...../phaseolus/index.html>.

VAN DER MAESEN is working on Leguminosae of Benin for the project Flora of Benin (1997-2002). Needs *Flemingia*, *Cajaniinae* and *Millettia* from Africa. Offers duplicates from Benin.

RECENT LEGUME LITERATURE

Ed. Note: Authors names in all capital letters are *Bean Bag* Readers. Their full names and addresses are listed in the *Bean Bag* Current Readers' Directory soon to be available on the www server of the Royal Botanic Gardens, Kew, United Kingdom. See "From the Editor" for the URL. All correspondence should be addressed directly to them.

ADEMA, F. 2000. Notes on Malesian Fabaceae (Leguminosae-Papilionoideae): 6. The rare genus *Burkilliodendron*. Gard. Bull. Singapore 52(1): 5-10. Icones, Anatomy and Morphology.

Akan, H. and S. Civelek. 2001. *Astragalus aytatchii* (Fabaceae), a new species from Anatolia, Turkey. Ann. Bot. Fenn. 38(3): 167-170. Icones, Maps, Anatomy and Morphology, Keys.

Aktoklu, E. 2001. Two new varieties and a new record in *Onobrychis* from Turkey. Turk. J. Bot. 25(5): 359-363. Icones, Anatomy and Morphology.

Ali, A.I. and D.D. SOKOLOFF. 2001. A new combination in *Pseudolotus* Rech. f. (Leguminosae: Loteae). Kew Bull. 56(3): 721-723. *Pseudolotus villosus* comb. nov.

Ali, H.B.M., A. Meister and I. Schubert. 2000. DNA content, rDNA loci, and DAPI bands reflect the phylogenetic distance between *Lathyrus* species. Genome 43(6): 1027-1032. Chromosome numbers, Molecular systematics.

Alm, T. and M. Piirainen. 2000. Krigsspredte arter i Sor-Varanger, Finnmark: knollerteknapp *Lathyrus linifolius*. (Polemochores in Sor-Varanger, Finnmark: *Lathyrus linifolius*.) Blyttia 58(3-4): 174-177. Icones, Maps.

Aniszews, T., M.H. Kupari and A.J. Leinonen. 2001. Seed number, seed size and seed diversity in Washington lupin (*Lupinus polyphyllus* Lindl.). Ann. Bot. (UK) 87(1): 77-82. Icones, Anatomy and Morphology, Reproductive biology.

Anon. 2001. Wild flowers of Kashmir: series 4, part 11. Sino-Himalayan Pl. Ass. Newsl. no.23: 21-22. Anatomy and Morphology.

ANULOV, O.V., N.I. Smirnova and E.V. Belyaeva. 2000. (Monosaccharide composition of seed coat structural polysaccharides in some species of legumes (chemotaxonomical aspect.) Byull. Glavn. Bot. Sada (Moscow) 179: 132-135. Chemotaxonomy.

Arevshatyan, I.G. 2001. (Revision of *Astragalus* (Fabaceae) species of Armenia: section Ornithopodium.) Bot. Zhurn. 86(4): 137-140. Icones, Keys.

Atahuachi, M., S. Arrazola and C. Antezana. 2001. Aporte al conocimiento del genero *Prosopis* (Mimosaceae) en Bolivia. (Approach to the botanical of the genus *Prosopis* in Bolivia.) Rev. Soc. Boliviana Bot. 3(1-2): 260-291. Icones, Maps, Anatomy and Morphology, Keys.

Athar, M. and J. Harding. 2000. Nodulating legumes from the Tahoe Basin, California. Sida, Contrib. Bot. 19(1): 205-210.

Aytac, Z. 2000. The genus *Ebenus* L (Leguminosae/Fabaceae) in Turkey. Karaca Arbor.Mag. 5(4): 145-171. Icones, Maps, Chromosome numbers, Anatomy and Morphology, Keys. English version pp.145-158

Aytac, Z., M. Ekici and L. Acik. 2001. A new species of *Astragalus* L. (Leguminosae) from south Anatolia. Bot. J. Linn. Soc. 136(1): 131-135. Icones, Maps, Anatomy and Morphology, Palynology, Molecular systematics. *Astragalus yilmazii* sp. nov.

Aytac, Z., F. Unal and M.N. Pinar. 2000. Morphological, palynological, and cytotaxonomical study of *Ebenus longipes* Boiss. & Bal. and *E. argentea* Siehe ex Bornm. (Leguminosae) from Turkey. Israel J. Pl. Sci. 48(4): 321-326. Icones, Chromosome numbers, Anatomy and Morphology, Palynology.

BANDYOPADHYAY, S. 2001. On the distribution range of *Bauhinia glauca* (Benth.) Benth. (Leguminosae: Caesalpinioideae) in India. J. Bombay Nat. Hist. Soc. 98(1): 149-150.

BANKS, H. and P. Gasson. 2000. Pollen morphology and wood anatomy of the *Crudia* group (Leguminosae, Caesalpinioideae, Detarieae). Bot. J. Linn. Soc. 134(1-2): 19-59. Icones, Anatomy and Morphology, Palynology, Keys.

- BANKS, H. and B.B. Klitgaard. 2001. Palynological contribution to the systematics of detarioid legumes (Leguminosae: Caesalpinioideae). In HERENDEEN, P.S. and A. BRUNEAU (eds.), Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.79-106. Palynology.
- BARKER, N.P., SCHRIRE, B.D. and J.H. Kim. 2001. Generic relationships in the tribe Indigoferae (Leguminosae: Papilionoideae) based on sequence data and morphology. In Herendeen, P.S. and Bruneau A. (eds.), Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.311-337. Anatomy and Morphology, Molecular systematics.
- Barrett, S. 2001. Lost and found: Drummond's grass and Cranbrook pea. *Danthonia* 10(1): 10.
- Bernatova, D. and J. Kliment. 2001. Zaujímavé floristické nálezky zo slovenských Karpat. (Interesting floristic findings from the Slovak Carpathians.) *Bull. Slov. Bot. Spolocn.* 23: 65-69.
- Bossi, F.S. and J.R. Davina. 2000. Cromosomas de cuatro especies de *Galactia* (Fabaceae). *Bonplandia* 10(1-4): 175-179. Chromosome numbers.
- Brouat, C., L. Gielly and D. MCKEY. 2001. Phylogenetic relationships in the genus *Leonardoxa* (Leguminosae: Caesalpinioideae) inferred from chloroplast trnL intron and trnL-trnF intergenic spacer sequences. *Amer. J. Bot.* 88(1): 143-149. Maps, Molecular systematics.
- Brullo, S., G. De-Marco and G. Siracusa. 2001. Taxonomical notes on the genus *Adenocarpus* DC. (Leguminosae) in Italy. *Bocconea* 13: 425-436. Icones, Maps, Anatomy and Morphology, Keys. 2 spp. nov.
- Brullo, S., R. Guarino and P. Minissale. 2000. Taxonomic and phytogeographical remarks on *Trifolium savianum* Guss., a misappreciated species of the Italian flora. *Bot. Jahrb.* 122(4): 469-480. Icones, Maps, Anatomy and Morphology.
- BRUNEAU, A., F.J. BRETELIER, J.J. WIERINGA, Gervais, G.Y.F and F. Forest. 2001. Phylogenetic relationships in tribes Macrolobieae and Detarieae as inferred from chloroplast trnL intron sequences. In HERENDEEN, P.S. and A. BRUNEAU (eds.), Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.121-149. Molecular systematics.
- BRUNEAU, A., F. Forest, P.S. HERENDEEN, B.B. KLITGAARD and G.P. LEWIS. 2001. Phylogenetic relationships in the Caesalpinioideae (Leguminosae) as inferred from chloroplast trnL intron sequences. *Syst. Bot.* 26(3): 487-514. Molecular systematics.
- Butcher, R. 2001. *Sphaerolobium benetectum* (Leguminosae: Mirbelieae), a new species from south-west Western Australia. *Nuytsia* 13(3): 449-455. Icones, Maps, Anatomy and Morphology, Keys.
- Butcher, R. and J.A. Chappill. 2001. *Sphaerolobium macranthum* (Leguminosae: Mirbelieae) complex revised. *Austral. Syst. Bot.* 14(2): 155-173. Icones, Maps, Anatomy and Morphology, Keys. *Sphaerolobium validum* sp. nov.
- CACCAVARI, M.A. 2000. Observaciones sobre la Morfología y estructura polínica de géneros del grupo "*Piptadenia*" (Mimosoideae, Leguminosae) de la América Tropical y Subtropical. Congress on Tropical and Subtropical Palynology. 9-14 Feb. 2001, La Habana, Cuba: 28.
- CACCAVARI, M.A. and E. Dome. 2000. An account of morphological and structural characterization of American Mimosoideae pollen. Part I, Tribe Acacieae, *Palynology*: 24 :231-248.
- Cajas, D. and C. Baeza. 2001. Anatomía del xilema secundario de las especies insulares Chilenas de *Sophora* L. (Fabaceae). (Secondary xylem of the Chilean insular species of *Sophora* L. (Fabaceae).) *Guyana, Bot.* 57(2): 181-184. Icones, Anatomy and Morphology.
- Calderon, J.S., F. Gomez-Garibay and C.L. Cespedes. 2001. Isoprenylated flavonoids from *Tephrosia tuitoensis*. *Biochem. Syst. Ecol.* 29(7): 763-764. Chemotaxonomy.
- Campbell-Young, G.J. and K. Balkwill. 2001. A new species of *Pearsonia* (Fabaceae) from dolomites in Northern Province, South Africa. *Nordic J. Bot.* 20(5): 547-555. Icones, Maps, Anatomy and Morphology, Keys. *Pearsonia callistoma* sp. nov.
- CARDENAS, L. and F. GARCIA. 2000. *Acacia guarensis* (Leguminosae-Mimosoideae), especie nueva del sur de Venezuela. *Ernstia* 10(4): 145-148. Icones, Anatomy and Morphology. *Acacia guarensis* sp. nov.

- Cardoso, M.B., M.T. Schifino-Wittmann and M.H. Bodanese-Zanettini. 2000. Taxonomic and evolutionary implications of intraspecific variability in chromosome numbers of species of *Leucaena* Benth. (Leguminosae). Bot. J. Linn. Soc. 134(4): 549-556. Chromosome numbers.
- Castroviejo, S. and others (eds.) 2000. Flora Iberica: plantas vasculares de la Peninsula Iberica e Islas Baleares. Vol. VII(II): Leguminosae (partim). Madrid: Real Jardin Botanico CSIC, 2000 xxii, 581-1119pp. Icones, Chromosome numbers, Anatomy and Morphology, Kcys. In memoriam Eugenio Sierra Rafols (11-v-1919/14-vii-1999) pp.xi-xiv.
- CHANDLER, G.T., R.J. Bayer and M.D. Crisp. 2001. A molecular phylogeny of the endemic Australian genus *Gastrolobium* (Fabaceae: Mirbelieae) and allied genera using chloroplast and nuclear markers. Amer. J. Bot. 88(9): 1675-1687. Maps, Molecular systematics.
- Chuakul, W. 2001. *Clitoria chanondii* (Leguminosae-Papilionoideae), a new species from Thailand. Kew Bull. 56(3): 737-740. Icones, Anatomy and Morphology.
- Cialdella, A.M. 2000. Flora fanerogamica Argentina: fasciculo 67. 128. Fabaceae, parte 9. Subfam. II. Mimosoideae, parte 3. Tribu VI. Mimoseae, parte A. Cordoba: Pro Flora, Conicet, 2000.10pp. Anatomy and Morphology, Keys.
- Clarke, H.D., S.R. Downie and D.S. SEIGLER. 2000. Implications of chloroplast DNA restriction site variation for systematics of *Acacia* (Fabaceae: Mimosoideae). Syst. Bot. 25(4): 618-632.
- Conte, L. and G. Cristofolini. 2000. Intraspecific diversity of *Cytisus emeriflorus* Reichenb. (Leguminosae), an endemic plant with disjunct distribution: evidence from isozyme data. Pl. Biosystems 134(3): 373-384. Maps.
- Coutinho, H.L.C.; Oliveira, V.M. and F.M.S. MOREIRA. 2000. Systematics of legume nodule nitrogen fixing bacteria. In: F.G. Priest and G. Goodfellow (eds.), Applied Microbial Systematics, Chapter 5, p107-134.
- Crespi, A.L., V. Morgado, A. Abreu, A. Pereira, J.A. Ribeiro, F. Amich and J. Paiva. 2000. Contribucion al conocimiento del genero *Lotus* L. en el norte de Portugal: I. Diversidad taxonomica y distribucion. Lagasalia 21(2): 279-287. Maps, Keys.
- CRISP, M.D., Gilmore, S. and B.E. VAN-WYK. 2001. Molecular phylogeny of the genistoid tribes of papilionoid legumes. In HERENDEEN, P.S. and A. BRUNEAU (eds.), Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.249-276. Molecular systematics.
- Da Silva, A.S.L and R.S. de Secco. 2000. *Mimosa dasilvae*, uma nova Mimosaceae da Amazonia Brasileira. (A new species of Mimosaceae from Brazilian Amazon, *Mimosa dasilvae*.) Acta Amazonica 30(3): 449-452. Icones, Anatomy and Morphology.
- Dangwal, L.R. and R.D. Gaur. 2000. Some rare and uncommon legumes from Garhwal Himalaya. J. Bombay Nat. Hist. Soc. 97(2): 309-311. Icones, Anatomy and Morphology.
- Davina, J.R. and A.I. Honfi. 2000. Los cromosomas de algunas especies lenosas. Bonplandia 10(1-4): 169-173. Chromosome numbers.
- DELGADO-SALINAS, A. 2000. New species of Mexican *Phaseolus* (Fabaceae). Syst. Bot. 25(3): 414-436. Icones, Chromosome numbers, Anatomy and Morphology, Palynology, Kcys. Many spp. nov.
- Devesa, J.A. 2000. Una nueva combinacion en *Trifolium*: *T. gemellum* var. *tastetii* (Pau) Devesa. Lagasalia 21(2): 365.
- DOYLE, J.J., J.A. CHAPPILL, C.D. Bailey and T. Kajita. 2001. Towards a comprehensive phylogeny of legumes: evidence from rbcL sequences and non-molecular data. In HERENDEEN, P.S. and A. BRUNEAU (eds.), Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.1-20.
- DOYLE, J.J., J.L. Doyle, A.H.D. Brown and B.E. Pfeil. 2000. Confirmation of shared and divergent genomes in the *Glycine tabacina* polyploid complex (Leguminosae) using histone H3-D sequences. Syst. Bot. 25(3): 437-448. Chromosome numbers.
- DREWES, S.I. and P.S. HOC. 2000. Morfologia y desarrollo de flores cleistogamas en *Macroptilium fraternum* (Fabaceae). (Morphology and development of cleistogamous flowers in *Macroptilium fraternum* (Fabaceae).) Kurtziana 28(2): 229-238. Icones, Anatomy and Morphology.

- Dupuy, D.J (ed.) 2001. The Legumes of Madagascar. Kew: Royal Botanic Gardens, Kew, 737 pp. Icones, Maps, Keys, Morphology, Illus, Gen.nov. & many spp. nov.
- Dusen, O.D. and H. Sumbul. 2000. *Glycyrrhiza asynmetrica* Hub.-Mor., egrimeyan, Fabaceae/Leguminosae (baklagiller). Karaca Arbor.Mag. 5(3): 137-140. Icones, Maps, Chromosome numbers, Anatomy and Morphology. English version pp.137-138
- Ebinger, J.E., D.S. SEIGLER and H.D. Clarke. 2000. Taxonomic revision of South American species of the genus *Acacia* subgenus *Acacia* (Fabaceae: Mimosoideae). Syst. Bot. 25(4): 588-617. Maps, Keys.
- Ebrahimzadeh, H., V. Niknam and A.A. MAASSOUMI. 2001. The sterols of *Astragalus* species from Iran: GLC separation and quantification. Biochem. Syst. Ecol. 29(4): 393-404.
- Edeoga, H.O., H.N. Ugbo and P.I. Osawe. 2000. Palynomorphology of species of *Commelina* L. and *Senna* Tourn. ex Mill. from Nigeria. New Bot. 25(1-4): 1-10. Icones, Palynology.
- Ekici, M. and Z. Aytac. 2001. *Astragalus dumanii* (Fabaceae), a new species from Anatolia, Turkey. Ann. Bot. Fenn. 38(3): 171-174. Icones, Maps, Anatomy and Morphology, Palynology.
- ESTRADA-C, A.E. and A. Martinez-M. 2000. Legumes from the central part of the State of Chihuahua, Mexico. Sida, Contrib. Bot. 19(2): 351-360. Maps.
- Etcheverry, A.V., S.M. Perez-de Bianchi, D.M. de Lopez and J.J. Protomastro. 2001. Floral biology of *Macroptilium erythroloma* (Benth.) Urban (Phaseoleae, Papilionoideae, Fabaceae). Beitr. Biol. Pfl. 71(3): 403-418. Icones, Anatomy and Morphology, Palynology, Reproductive biology.
- Flores-Franco, G. and M. SOUSA-S. 2000. Especie nueva de *Chamaecrista* (Leguminosae, Caesalpinioideae) de Mesoamerica. An. Inst. Biol. Univ. Nac. Auton. Mex., Bot. 71(1): 19-23. Icones, Anatomy and Morphology. *Chamaecrista molinae* sp. nov.
- FORTUNATO, R.H. 2001. Systematic relationships in Rhynchosia (Cajaninae-Phaseoleae-Papilionoideae-Fabaceae) from the neotropics. In HERENDEEN, P.S. and A. BRUNEAU (eds.), Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp. 339-354. Maps.
- FUENTES, E.F., and A.M. PLANCHUELO. 2000. Wild lupins as pioneers of riverside sand banks. In van Santem, E., M. Wink, S. Weissmann & P. R  emer (eds.), Lupin, an Ancient crop for the New Millenium: 316-319. Auburn Univ. USA. ISBN 0-86476-123-6. Ecology.
- Gamerro, J.C. and R.H. FORTUNATO. 2001. Morfologia del polen de las especies Argentinas de *Bauhinia* (Cercideae, Caesalpinioideae, Fabiaceae). Ann. Missouri Bot. Gard. 88(1): 144-158. Icones, Palynology.
- GARCIA, R. and M. Mejia. 2000. Una especie de *Acacia* (Mimosaceae) nueva para la ciencia. Moscosoa 11: 7-10. Icones, Anatomy and Morphology. *Acacia oviedoensis* sp. nov.
- GASSON, P. 2001. Does wood anatomy support tribal and generic classification in papilionoid Leguminosae? In HERENDEEN, P.S. and A. BRUNEAU (eds.), Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.201-215. Icones, Anatomy and Morphology.
- GASSON, P. and E.J. Wray. 2001. Wood anatomy of *Cyathostegia mathewsii* (Swartzieae, Papilionoideae, Leguminosae). IAWA J. 22(2): 193-199. Icones, Anatomy and Morphology.
- Germishuizen, G., W. Coetzer and G. Condry. 2001. *Caesalpinia rostrata*. Flow. Pl. Afr. 57: 70-74. Icones, Maps, Anatomy and Morphology.
- Gesti, J. and E. Fabregas. 2000. *Amorpha fruticosa* L. (Papilionaceae) al Baix Emporda. (*Amorpha fruticosa* L. (Papilionaceae) in the Baix Emporda (NE Catalonia).) But. Inst. Catalana Hist. Nat. 68: 90-92, 95-96. Icones, Anatomy and Morphology.
- Ghahremani-nejad, F. 2001. Notes on *Astragalus* sect. *Corethrui* (Fabaceae), including a new combination. Ann. Bot. Fenn. 38(1): 47-49. Anatomy and Morphology. *Astragalus scoparius* subsp. *intarrensii* comb. nov.

- GOMEZ-SOSA, E. 2000. Las especies Argentinas de *Crotalaria* (Leguminosae-Crotalarieae): novedades, descripciones y clave. (Species of *Crotalaria* (Leguminosae-Crotalarieae) from Argentina: new reports, descriptions and key.) Gayana, Bot. 57(1): 67-87. Icons, Chromosome numbers, Anatomy and Morphology, Keys.
- Gomiz-Garcia, F. 2000. Notas sobre flora de Marruecos, 3. An. Jard. Bot. Madrid 58(2): 364-365. Icons, Anatomy and Morphology. *Rhodanthemum kesticum* sp. nov.
- Grankina, V.P. 2001. (Generis *Glycyrrhiza* L. (Fabaceae) species novae ex Asia Centrali.) Novosti Sist. Vyssh. Rast. 33: 145-150. Anatomy and Morphology.
- Gurusamy, C. and A.K. Bal. 2000. Effect of seed abortion on the pattern of seed formation in *Lathyrus maritimus* and *L. sativus* (Fabaceae). Phytion (Austria) 40(2): 223-238. Icons.
- Hamzehee, B. 2000. Some new and noteworthy plant records from Iran. Iranian J. Bot. 8(2): 271-277. Icons, Anatomy and Morphology.
- HERENDEEN, P.S. 2001. Structural evolution in the Caesalpinioideae (Leguminosae). In Herendeen P.S. and Bruneau, A. (eds.), Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.45-64. Icons, Anatomy and Morphology.
- HERENDEEN, P.S. and A. BRUNEAU (eds.). 2000. Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 x, 363pp. Icons, Anatomy and Morphology, Palynology, Reproductive biology, Molecular systematics.
- Hervas-Serrano, J.L., A.M. Fernandez-Ocana, J.M. Espinosa-Gento and C. Fernandez-Lopez. 2000. Algunas Labiadas y Leguminosas del Alto Guadalquivir (sur de la peninsula Iberica) hasta 1.999. Blancoana no.17: 89-90.
- Holman, J.E. and J. Playford. 2000. Molecular and morphological variation in the *Senna artemisioides* complex. Austral. J. Bot. 48(5): 569-579. Maps, Chromosome numbers, Anatomy and Morphology, Molecular systematics.
- Hopkins, H.C.F. 2000. *Parkia paya* (Leguminosae: Mimosoideae), a new species from swamp forest and notes on variation in *Parkia speciosa* sensu lato in Malesia. Kew Bull. 55(1): 123-132. Icons, Maps, Anatomy and Morphology, Keys.
- Hsu, Y.C. and T.C. Huang. 2001. A palynological study of the tribe Millettieae (Fabaceae) in Taiwan. Taiwanica 46(1): 21-39. Icons, Anatomy and Morphology, Palynology.
- Hurter, P.J.H. and E. van Wyk. 2001. First distribution record for *Brachystegia spiciformis* (Caesalpinioideae) in South Africa. Bothalia 31(1): 43-44. Maps.
- Hu, J.M. 2001. Phylogenetic relationships of the tribe Millettieae and allies: the current status. In Herendeen P.S. and A. Bruneau (eds.), Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.299-310. Molecular systematics.
- IRELAND, H., R.T. PENNINGTON and J. Preston. 2001. Molecular systematics of the Swartzieae. In Herendeen P.S. and A. Bruneau (eds.), Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.217-231. Molecular systematics.
- JAASKA, V. 2001. Isoenzyme diversity and phylogenetic relationships among the American beans of the genus *Vigna* Savi (Fabaceae). Biochem. Syst. Ecol. 29(11): 1153-1173. Chemotaxonomy.
- Jawad, J.T., D.S. SEIGLER and J.E. Ebinger. 2000. A systematic treatment of *Acacia coulteri* (Fabaceae, Mimosoideae) and similar species in the New World. Ann. Missouri Bot. Gard. 87(4): 528-548. Maps, Anatomy and Morphology, Keys. *Acacia durangensis* comb. nov.
- Jobson, P.C. and P.H. WESTON. 2001. *Dillwynia rupestris* (Fabaceae: Mirbelieae), a new species from the New England Tableland of New South Wales. Telopea 9(2): 323-327. Icons, Anatomy and Morphology.
- JOHNSON, M.B. 2001. Name changes of legumes used in southwest landscapes. Desert Pl. 17(1): 30-31. Icons.
- Kajita, T. and H. OHASHI. 2001. Intraspecific restriction site variation in the chloroplast DNA of *Desmodium podocarpum* DC. (Leguminosae-Papilionoideae). J. Jap. Bot. 76(2): 77-83. Icons, Molecular systematics.

- Kajita, T., H. OHASHI, Y. Tateishi, C.D. Bailey and J.J. DOYLE. 2001. RbcL and legume phylogeny, with particular reference to Phaseoleae, Millettieae, and allies. *Syst. Bot.* 26(3): 515-536. Molecular systematics.
- Katsuyama, T., A. Sasaki and M. Igarashi. 2001. *Vicia lathyroides* L. (Leguminosae), naturalized in Japan. *J. Jap. Bot.* 76(3): 173-174. Icones.
- KEIGHERY, G.J. 2001. A new subspecies of *Isotropis cuneifolia* (Fabaceae). *Nuytsia* 13(3): 471-474. Icones, Anatomy and Morphology. *Isotropis cuneifolia* subsp. *glabra* subsp. nov.
- Kholina, A.B., O.G. Koren and Y.N. Zhuravlev. 2000. Allozyme variation in *Oxytropis retusa* Matsum. from the Kuril Archipelago. *Nat. Hist. Res. (Chiba) special issue no.7*: 15-20. Maps.
- KLITGAARD, B.B., L.P. de QUEIROZ and G.P. LEWIS. 2000. A remarkable new species of *Pterocarpus* (Leguminosae: Papilionoideae: Dalbergieae) from Bahia, Brazil. *Kew Bull.* 55(4): 989-992 (2000) - illus.
- Knyazev, M.S. 2001. Zametki po sistematike i khorologii vidov roda *Oxytropis* (Fabaceae) na Urale: 2. Vidy rodstva *Oxytropis ambigua*. (Systematic and chorological notes on the species of the genus *Oxytropis* (Fabaceae) in the Urals: 2. Species related to *Oxytropis ambigua*.) *Bot. Zhurn.* 86(1): 126-134. Icones, Maps, Chromosome numbers, Anatomy and Morphology. 2 spp. nov.; 1 subsp. nov.; 1 var. nov.
- Kodala, P.G. 2001. *Acacia atrox* (Fabaceae: Mimosoideae), a new rare species from the North Western Slopes, New South Wales. *Telopea* 9(2): 415-419. Icones, Anatomy and Morphology.
- Kodala, P.G. and M.D. TINDALE. 2001. *Acacia dealbata* subsp. *subalpina* (Fabaceae: Mimosoideae), a new subspecies from south-eastern Australia. *Telopea* 9(2): 319-322. Anatomy and Morphology, Keys.
- Kodala, P.G., M.D. TINDALE and D.A. Keith. 2001. *Acacia stellaticeps* (Fabaceae: Mimosoideae), a new species from the Pilbara, Western Australia, to Tanami, Northern Territory. *Nuytsia* 13(3): 483-486. Icones, Anatomy and Morphology.
- Kotresha, K. and Y.N. Seetharam. 2000. Epidermal micromorphology of some species of *Cassia* L. (Caesalpinaceae). *Phytomorphology* 50(3-4): 229-237. Icones, Anatomy and Morphology, Keys.
- KUMAR, S. 2000. Synoptic account of Leguminosae (nom. alt. Fabaceae) of Haryana. In Gupta, B.K. (ed.), Higher plants of Indian subcontinent, vol. IX. Dehra Dun: Bishen Singh Mahendra Pal Singh, 2000 pp.169-200. (Indian Journal of Forestry Additional series; 11.)
- Laguna-Lumbreras, E. 2001. Sobre el frijol flamenco o careto, *Lablab purpureus* L. Sweet, en tierras Valencianas. *Flora Montiberica* 17: 12-20. Anatomy and Morphology.
- LARSEN, K. and S.S. LARSEN. 2000. *Bauhinia strychnoidea* Prain (Leguminosae-Caesalpinioideae), a new record from Thailand. *Thai Forest Bull., Bot.* no.28: 33-37. Icones, Anatomy and Morphology.
- Lavia, G.I. 2001. Chromosomal characterization of germplasm of wild species of *Arachis* L. belonging to sections *Trirectoides*, *Erectoides* and *Procumbentes*. *Caryologia* 54(2): 115-119. Chromosome numbers.
- LAVIN, M., R.T. PENNINGTON, B.B. KLITGAARD, J.I. SPRENT, de-Lima, H.C. and P.E. GASSON. 2001. The dalbergioid legumes (Fabaceae): delimitation of a pantropical monophyletic clade. *Amer. J. Bot.* 88(3): 503-533. Icones, Anatomy and Morphology, Molecular systematics.
- LAVIN, M., M. Thulin, J.N. Labat and R.T. PENNINGTON. 2000. Africa, the odd man out: molecular biogeography of dalbergioid legumes (Fabaceae) suggests otherwise. *Syst. Bot.* 25(3): 449-467.
- LEWIS, G.P., B.B. Simpson and J.L. Neff. 2001. Progress in understanding the reproductive biology of the Caesalpinioideae (Leguminosae). In Herendeen P.S. and A. Bruneau (eds.), *Advances in legume systematics: part 9*. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.65-78. Reproductive biology.
- Linares, J. 2001. Nuevas especies del genero *Ateleia* (Leguminosae: Papilionoideae) de Mexico y Centroamerica. (New Mexican and Central American species of *Ateleia* (Leguminosae: Papilionoideae).) *An. Inst. Biol. Univ. Nac. Auton. Mex. Bot.* 72(1): 85-114. Icones, Anatomy and Morphology, Keys. Many spp. nov.

- Liu, C.C. and T.C. Huang. 2001. Morphological evidences for hybrid of *Dumasia* (Fabaceae) in Taiwan. *Taiwania* 46(1): 1-12. Icones, Anatomy and Morphology, Palynology.
- Liu, L.F., G.Z. Qian and Y.Z. Zhang. 2000. (The taxonomic study on *Oxytropis moellendorffii* Bunge (Legumen).) *Bull. Bot. Res. (China)* 20(4): 369-371.
- Liu, Y.B. and Y.Z. Zhao. 2000. (A cladistic taxonomic method on studying the systemic evolution of the genus *Vicia* L. in the Mongolian Plateau.) *Acta Sci. Nat. Univ. Neimongol* 31(3): 293-299. Keys.
- Li, S.F., B. Li, Z.Y. Chang and L.R. Xu. 2000. (A cytological study of *Astragalus dshimensis* Gontsh. and *A. hoantchy* Franch. with a taxonomic revision.) *Acta Bot. Bor. Occid. Sin.* 20(3): 415-420. Maps, Chromosome numbers.
- Li, S.J., D.X. Zhang and Z.Y. Chen. 2001. (A species of *Caesalpinia* L. (Leguminosae) new to China.) *Guihaia* 21(2): 106. Chromosome numbers.
- Lopez, J., J.A. Devesa, A. Ortega-Olivencia and T. Ruiz. 2000. Production and morphology of fruit and seeds in Genisteae (Fabaceae) of south-west Spain. *Bot. J. Linn. Soc.* 132(2): 97-120. Icones, Anatomy and Morphology, Keys.
- Losano, M.A., N. Dottori and M.T. Cosa. 2000. Secreciones intravasculares de sustancias gomosas en *Cercidium praecox* (Fabaceae). *An. Inst. Biol. Univ. Nac. Auton. Mex., Bot.* 71(1): 1-9. Icones, Anatomy and Morphology.
- LUCKOW, M., P.J. White and A. BRUNEAU. 2001. Relationships among the basal genera of mimosoid legumes. In Herendeen P.S. and A. Bruneau (eds.), *Advances in legume systematics: part 9*. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.165-180. Molecular systematics.
- Mackinder, B.A. 2001. The monophyly and taxonomic position of *Berlinia* Hook. f. (Leguminosae-Caesalpinioideae): a preliminary study. In Herendeen P.S. and A. Bruneau (eds.), *Advances in legume systematics: part 9*. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.151-163. Anatomy and Morphology, Molecular systematics.
- Malaviya, D.R. and R.S. Shukla. 2000. Evolutionary trend in *Lens culinaris* and allied species: a cytological evidence. *Cytologia (Japan)* 65(3): 305-311. Chromosome numbers.
- Marino-S, N.M. 2000. Caracterizacion morfoanatomica de la cubierta seminal en *Canavalia ensiformis* (L.) DC. cv. Tovar. *Ernstia* 10(3): 85-93. Anatomy and Morphology.
- Marticorena, C. 2001. Nuevos nombres y nuevas combinaciones en la flora de Chile. (New names and combinations for the Chilean flora.) *Guyana, Bot.* 57(2): 191-192. Many comb. nov.
- Martinez, J. and M. Mayor. 2000. La distribucion de *Vicia pubescens* (DC.) Link e *Hippocrepis commutata* Pau en la Cornisa Cantabrica. *Bot. Cienc. Nat. I.D.E.A.* no.46: 281-287. Maps.
- Merino, E. F., A.M. PLANCHUELO and M. Wink. 2000. Phylogenetic analysis of *Lupinus*. In van Santem, E., M. Wink, S. Weissmann & P. R  emer (eds). *Lupin, an Ancient crop for the New Millenium*: 287-290. Auburn Univ. USA. ISBN 0-86476-123-6. Molecular systematics.
- Merino, E.F., E. FUENTES and A.M. PLANCHUELO. 2000. Phytochemical characterization of lupin species of South America. In van Santem, E., M. Wink, S. Weissmann & P. R  emer (eds). *Lupin, an Ancient Crop for the New Millennium*: 291-293. Auburn Univ. USA. ISBN 0-86476-123-6. Chemotaxonomy.
- Miguel, D.L. and F.M.S. MOREIRA. 2001. Influence of medium and peat pH on the behaviour of *Bradyrhizobium* strains. *Revista Brasileira de Ci  ncia do Solo*, 25: 873-883.
- Mckey, D.B. 2000. *Leonardoxa africana* (Leguminosae: Caesalpinioideae): a complex of mostly allopatric subspecies. *Adansonia* 22(1): 71-109. Icones, Maps, Anatomy and Morphology, Keys. 3 subsp. nov.
- MILLER, J.T. and R.J. Bayer. 2001. Molecular phylogenetics of *Acacia* (Fabaceae: Mimosoideae) based on the chloroplast trnK/matK and nuclear Histone H3-D DNA sequences. In Herendeen P.S. and A. Bruneau (eds.), *Advances in legume systematics: part 9*. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.181-200. Molecular systematics.

- Mollard, F.P.O., P.S. HOC and R.A. PALACIOS. 2000. *Prosopis abbreviata* (Mimosaceae) y su presunto origen híbrido. (*Prosopis abbreviata* (Mimosaceae) and its putative hybrid origin.) Bol. Soc. Argent. Bot. 35(3-4): 305-313. Icones, Palynology.
- MOREIRA, F.M.S. 2000. Biodiversity of *Rhizobia* from a wide range of forest Leguminosae species in Brazil. In: Nitrogen fixation: from molecules to crop productivity. Pedrosa, F.O.; Hungria, M.; Yates, M.G. and W.E. Newton (eds.). Kluwer Academic Publishers, pp.181-182.
- MOREIRA, F.M.S.; Carvalho, Y.; Gonçalves, M.; Haukka, K.; Young, P.J.; Faria, S.M.; Franco, A.A.; Cruz, L.M. and F.O. Pedrosa. 2000. *Azorhizobium johanense* sp.nov. and *Sesbania virgata* (Caz.) Pers.: a highly specific symbiosis. In: Nitrogen fixation: from molecules to crop productivity. Pedrosa, F.O.; Hungria, M.; Yates, M.G.; Newton, W.E. (eds.). Kluwer Academic Publishers, pp.197.
- MOREIRA, F.M.S. and E. Pereira. 2001. Microsymbionts: *Rhizobia*. In: M. Swift and D. Bignell (eds.): Standard methods for assessment of soil biodiversity and land-use practice. Bogor:ASB Lecture Note 6B/ICRAF, 34 pp.
- MURPHY, D.J., F. Udovicic and P.Y. Ladiges. 2000. Phylogenetic analysis of Australian *Acacia* Leguminosae: Mimosoideae by using sequence variations of an intron and two intergenic spacers of chloroplast DNA. Austral. Syst. Bot 13(5): 745-754. Molecular systematics.
- Murugesan, R., N. Chandra, S.T. Santhiya, P.M. Gopinath and K.M. Marimuthu. 2001. Nucleolar behaviour in a few legumes. Caryologia 54(1): 17-21.
- Mutlu, B. 2001. *Chesneya elegans* Fomin Fabaceae/Leguminosae. Karaca Arbor. Mag. 6(1): 31-34. Icones, Maps, Anatomy and Morphology. English version pp.31-33.
- Nagesh, K., G.K. Gaud, M.P. Raj and P.S. Rao. 2001. Tricotyledony in *Prosopis cineraria* (L.) Druce. (Mimosaceae). Indian Forester 127(5): 591-592.
- Nagesh, K., C.S. Reddy and M.R. Bhanja. 2001. Tricotyledony in *Acacia mellifera* (Vahl) Benth. (Mimosaceae). Indian J. Forest. 23(4): 440-441. Icones, Anatomy and Morphology.
- Nazneen-Parveen, S., K. Sri-Rama-Murthy and T. Pullaiah. 2000. Leaf epidermal characters in *Crotalaria* species (Papilionoideae) from Eastern Ghats. Phytomorphology 50(2): 205-212. Icones, Anatomy and Morphology, Keys.
- Ohba, H., S. AKIYAMA and S.K. Wu. 2000. Five new species of *Oxytropis* from the Kunlun and the Hoh Xil Mountains, NW China (Fabaceae). Bull. Nation. Sci. Mus., B (Tokyo) 26(2): 43-59. Icones, Anatomy and Morphology.
- Orchard, A.E. and A.J.G. Wilson (eds.). 2001. Flora of Australia: volume 11A. Mimosaceae, *Acacia* part 1. Canberra, A.C.T.: Australian Biological Resources Study, 2001 xx, 673pp. Icones, Maps, Anatomy and Morphology, Keys. Many new taxa.
- ORTIZ, S., I. Pulgar and I. Iglesias. 2001. A new species of *Cytisus* Desf. (Fabaceae) from islands off the west coast of Galicia (north-west Iberian Peninsula). Bot. J. Linn. Soc. 136(3): 339-344. Icones, Anatomy and Morphology, Palynology. *Cytisus insularis* sp. nov.
- Pakravan, M., N. Jalilian and M. Neamati. 2000. Flora of Iran: no. 33. Papilionaceae (Vicieae). Tehran: Research Institute of Forests and Rangelands, 2000. 176pp. Icones, Maps, Anatomy and Morphology, Keys.
- PALACIOS, R.A. and P.S. HOC. 2001. Three new species of *Prosopidastrum* (Mimosaceae) from Argentina. Novon 11(1): 79-87. Icones, Maps, Anatomy and Morphology, Palynology, Keys.
- Pardo, C., H. Tahiri, P. Cubas and F.E. el Alaoui-Faris. 2000. Pollen morphology in *Cytisus* (Papilionoideae, Leguminosae) from Morocco and the Iberian Peninsula. Grana 39(4): 159-168. Icones, Palynology.
- PASQUET, R.S. 2001. Notes on the genus *Vigna* (Leguminosae-Papilionoideae). Kew Bull. 56(1): 223-227. 1 comb. nov.; 1 var. nov.
- PAVLOVA, D.K. and V.I. Manova. 2000. Pollen morphology of the genera *Onobrychis* and *Hedysarum* (Hedysareae, Fabaceae) in Bulgaria. Ann. Bot. Fenn. 37(3): 207-217. Icones, Palynology.
- PAVLOVA, M.E. 2000. Seed morphology of *Bauhinia* L. (Fabaceae-Caesalpinioideae) applicable to classification. Dostignia I perspektivi v oblasti tropicheskogo zemledelia I givotnovodstva. (Achievements and perspectives in the sphere of tropical animal husbandry), Russian Univ. People's Friendship: 77. Russian with English summary.

- Peguero, B., A. Veloz, J. Salazar and R. Bastardo. 2000. Notas sobre la flora de la Isla Espanola: 7. Moscosoa 11: 107-112. Icones, Anatomy and Morphology.
- Penalzoa-A-del P-de, S. 2000. Citogenetica das especies silvestres do genero *Arachis* (Leguminosae). In Topicos atuais em botanica: palestras convidadas do 51st Congresso Nacional de Botanica, 23-29 de julho de 2000. Brasilia: Embrapa Recursos Geneticos e Biotecnologia, 2000 pp.45-49. Chromosome numbers.
- PENNINGTON, R.T. and B. Gemeinholzer. 2000. Cryptic clades, fruit wall morphology and biology of *Andira* (Leguminosae: Papilionoideae). Bot. J. Linn. Soc. 134(1-2): 267-286. Icones, Anatomy and Morphology.
- PENNINGTON, R.T., B.B. KLITGAARD, H. IRELAND and M. LAVIN. 2001. New insights into floral evolution and basal Papilionoideae from molecular phylogenies. In Herendeen P.S. and Bruneau A. (eds.), Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.233-248. Icones, Maps, Anatomy and Morphology, Molecular systematics.
- PENNINGTON, R.T., M. LAVIN, H. IRELAND, B. KLITGAARD, J. Preston and J.M. Hu. 2001. Phylogenetic relationships of basal papilionoid legumes based upon sequences of the chloroplast trnL intron. Syst. Bot. 26(3): 537-556. Molecular systematics.
- PENNINGTON, R.T., G. LEWIS and M. Marsh. 2000. Plant portraits: 399. *Andira inermis* subsp. *inermis*. Leguminosae-Papilionoideae. Curtis's Bot. Mag. 17(4): 188-194. Icones, Anatomy and Morphology.
- Phan-Ke, Loc and J.E. VIDAL. 2001. Flore du Cambodge, du Laos et du Vietnam: 30. Leguminosae-Papilionoideae Millettieae. Paris: Museum national d'histoire naturelle, 2001. 191pp. Icones, Anatomy and Morphology, Keys. Many new taxa.
- PIERGIOVANNI, A.R., D. Cerbino & M. Brandi. 2000. The common bean populations from Basilicata (Southern Italy). An evaluation of their variation. Genet. Res. Crop. Evol. 47: 489-495.
- PIERGIOVANNI, A.R., D. Cerbino & C. Della Gatta. 2000. Diversity in seed quality traits of common bean (*Phaseolus vulgaris*) populations from Basilicata (Southern Italy). Plant Breeding 119: 513-516.
- PIERGIOVANNI, A.R., G. Tarant & D. Pignone. 2000. Diversity among common bean populations from the Abruzzo region (Central Italy): a preliminary inquiry. Genet. Res. Crop. Evol. 47: 467-470.
- Pina, F.J. 2000. Novedades corologicas para la flora de Andalucia occidental. Lagasalia 21(2): 349-351.
- Pistelli, L., A. Bertoli, I. Giachi, I. Morelli, P. Rubiolo and C. Bicchi. 2001. Quinolizidine alkaloids from *Genista ephedroides*. Biochem. Syst. Ecol. 29(2): 137-141. Chemotaxonomy.
- Pitarch, R. and E. Sanchis-Duato. 2001. *Coronilla minima* L. subsp. *vigoi*, subespecie nueva para la flora iberica. Flora Montiberica 17: 21-23. Icones, Anatomy and Morphology.
- PLANCHUELO, A.M. 2000. Endangered species of wild lupins in South America. In van Santem, E., M. Wink, S. Weissmann & P. Röemer (eds). Lupin, an Ancient crop for the New Millenium: 34-37. Auburn Univ. USA. ISBN 0-86476-123-6. Ecology. Endangered species.
- PODLECH, D. 2001. Contributions to the knowledge of the genus *Astragalus* L. (Leguminosae): 7-10. Sendtnera 7: 163-201. Anatomy and Morphology, Keys. Many new taxa.
- PODLECH, D. and A.A. Maassoumi. 2001. New species of *Astragalus* L. sect. *Hymenostegis* (Fabaceae) from Iran. Sendtnera 7: 203-209. Anatomy and Morphology. Many spp. nov.
- Polatschek, A. 2000. Flora von Nordtirol, Osttirol und Vorarlberg: Bd 3. Samenpflanzen: Fabaceae bis Rosaceae. Kartenteil. Innsbruck: Tiroler Landesmuseum Ferdinandeum, 1354pp. Maps.
- PONOMARENKO, S. 2000. The results of research of leguminous seed microscopic structure (family Fabaceae) and their evolution. Dostignia I perspektivi v oblasti tropicheskogo zemledelia I givotnovodstva. (Achievements and perspectives in the sphere of tropical animal husbandry), Russian Univ. People's Friendship: 71. Russian with English summary.
- Pope, G.V. ed. 2000. Flora Zambesiaca: Mozambique, Malawi, Zambia, Zimbabwe, Botswana. Volume three, part six. Kew: Royal Botanic Gardens, Kew, for the Flora Zambesiaca Managing Committee, 2000 vi, 175p. Icones, Anatomy and Morphology. Keys. 2 spp. nov.; 2 comb. nov.

- Pullaiah, T. and K. Sri-Ramamurthy. 2001. Flora of Eastern Ghats: hill ranges of south east India. Vol. 2, Leguminosae (Fabaceae). New Delhi: Regency, 2001 viii, 12pp. of plates, 387pp. Icones, Anatomy and Morphology, Keys.
- Qian, Y.Y. 2001. *Millettia pendula* Benth. (Papilionaceae), a new record to China. Acta Phytotax. Sin. 39(5): 464-465. Icones, Anatomy and Morphology.
- Rabelo, L.A., F.M. de Agra, E.V.L. da Cunha, M.S. DA SILVA and J.M. Barbosa-Filho. 2001. Homohesperetin and phaseollidin from *Erythrina velutina*. Biochem. Syst. Ecol. 29(5): 543-544. Chemotaxonomy.
- Ramirez-Medina-ME-del, C. and J.J. Reynoso-Duenas. 2000. Riqueza y distribucion de Caesalpinaceae en el occidente de Mexico. Bol. Inst. Bot. (Guadalajara) 7(1-3): 1-38. Maps.
- Randell, B.R. 2000. *Cassia* and *Senna*: a review of the groups of plants formerly known as Cassia. Austral. Pl. 20(162): 238-249. Icones, Anatomy and Morphology.
- Reddy, C.S., M.R. Bhanja and V.S. Raju. 2000. *Cassia uniflora* Miller: a new record for Andhra Pradesh, India. Indian J. Forest. 23(3): 324-325. Anatomy and Morphology.
- Richardson, J.E., R.T. PENNINGTON, T.D. Pennington and P.M. Hollingsworth. 2001. Rapid diversification of a species-rich genus of neotropical rain forest trees. Science 293(5538): 2242-2245. Maps.
- Rich, E. 2000. More on lady's slipper in Hampshire. BSBI News no.84: 28.
- Rich, T.C.G. 2001. What is *Anthyllis vulneraria* L. subsp. *corbieri* (Salmon and Travis) Cullen (Fabaceae)? Watsonia 23(4): 469-480. Maps.
- RICO-ARCE, L. and H. BANKS. 2001. A preliminary survey of pollen and other morphological characters in neotropical *Acacia* subgenus* *Aculeiferum* (Leguminosae: Mimosoideae). Bot. J. Linn. Soc. 135(3): 263-270. Anatomy and Morphology, Palynology.
- RICO-ARCE, L. 2000. El genero *Acacia* (Leguminosae, Mimosoideae) en el Estado de Oaxaca, Mexico. (The genus *Acacia* (Leguminosae, Mimosoideae) in the state of Oaxaca, Mexico.) An. Jard. Bot. Madrid 58(2): 251-302. Icones, Maps, Anatomy and Morphology, Keys. 2 comb. nov.
- RICO-ARCE, L., M.S. SOUSA and S. FUENTES. 2000. Guinetia: a new genus in the tribe Ingeae (Leguminosae: Mimosoideae) from Mexico. Kew Bull. 54(4): 975-981. Icones, Anatomy and Morphology, Palynology. Guinetia tehuantepecensis sp. nov.
- RICO-ARCE, L. 2000. A new name for *Albizia elegans* (Ducke) L. Rico (Leguminosae: Mimosoideae). Kew Bull. 55(2): 404.
- Rizzi-Longo, L., L. Feoli-Chiapella, T. Cusma-Velari and V. Kosovel. 2000. Pollen cards of the *Genista radiata* group (Genisteae, Fabaceae). Stud. Geobot. 19: 39-48. Icones, Anatomy and Morphology, Palynology.
- Rodriguez-Marzal, J.L. 2000. Tres plantas nuevas para la flora onubense. Lagasalia 21(2): 336-338.
- ROSS, J.H. 2001. Two new endemic Australian genera in the tribe Brongniartieae (Fabaceae) to accommodate two species formerly included in *Templetonia* R. Br. Muelleria 15: 7-14. Chemotaxonomy, Keys, Molecular systematics.
- Rozan, P., Y.H. Kuo and F. Lambein. 2001. Amino acids in seeds and seedlings of the genus *Lens*. Phytochemistry 58(2): 281-289. Chemotaxonomy.
- Sahai, K. 2001. Anatomical variability in seed coat of some *Cassia* L. (Caesalpinioideae) species with taxonomic significance. Taiwania 46(2): 158-166. Icones, Anatomy and Morphology.
- Sahai, K. and S.M. Singh. 2001. Macro- and micro-morphological features of seeds in six genera of the tribe Phaseoleae (subfamily Papilionoideae). Phytomorphology 51(2): 191-197. Icones, Anatomy and Morphology, Reproductive biology.
- Saidman, B.O., C.F. Bessega, L.I. Ferreyra, N. Julio and J.C. Vilardi. 2000. The use of genetic markers to assess population structure and relationships among species of the genus *Prosopis* (Leguminosae). Bol. Soc. Argent. Bot. 35(3-4): 315-324.

- Sanchir, C.h. 2000. Dva novykh vida roda *Caragana* Lam. (Fabaceae) iz Kazakhstana. (Generis *Caragana* Lam. (Fabaceae) species duae novae e Kazachstania.) Novosti Sist. Vyssh. Rast. 32: 72-76. 2 spp. nov.; 2 var. nov.
- SARAVANAKUMAR, P. 2000 Cluster analysis of some wild relatives of pulses. Proc. 12th Annual Sess. Inst. Biol. Sept. 2000. Sri Lanka.
- SARAVANAKUMAR, P. 2000. Establishment of potential relationships between two grain legumes and their tentative wild relative. Annual Research Sessions. October 2000. Univ. Paradenrya.
- Sartori, A.L.B. and M.G. Azevedo-Tozzi. 2001. *Myrocarpus emarginatus* (Leguminosae, Papilionoideae, Sophoreae), a new species from Bolivia. Novon 11(3): 353-355. Icones, Anatomy and Morphology.
- Sawkins, M.C., B.L. MAASS, B.C. Pengelly, H.J. Newbury, B.V. Ford-Lloyd, N. MAXTED and R. Smith. 2001. Geographical patterns of genetic variation in two species of *Stylosanthes* Sw. using amplified fragment length polymorphism. Molec. Ecol. 10(8): 1947-1958. Chromosome numbers, Molecular systematics.
- Sa, R. and C.J. Chen. 2000. Surface features of seeds in *Thermopsis* (Leguminosae). Acta Phytotax. Sin. 38(6): 582-587. Anatomy and Morphology, Reproductive biology. 1pp. illus. at end of issue.
- Sa, R., C.J. Chen and P.C. Li. 2000. The phytogeographical studies of *Thermopsis* (Fabaceae). Acta Phytotax. Sin. 38(2): 148-166. Maps.
- Sa, R. and C.R. Chen. 2001. Examination of the type and related species in *Thermopsis* R. Br. Aiton et W. T. Aiton (Leguminosae). Acta Bot. Yunnanica 23(1): 41-42.
- Schiemann, K., T. Tyler and B. Widen. 2000. Allozyme diversity in relation to geographic distribution and population size in *Lathyrus vernus* (L.) Bernh. (Fabaceae). Pl. Syst. Evol. 225(1-4): 119-132. Maps.
- Seijo, G. 2000. Numeros cromosomicos de especies de *Mimosa* (Leguminosae) de Paraguay. Bonplandia 10(1-4): 163-167. Chromosome numbers.
- Seijo, G. and A. Fernandez. 2001. Chromosome numbers of some southernmost species of *Mimosa* L. (Leguminosae). Cytologia (Japan) 66(1): 19-23. Maps, Chromosome numbers.
- Semmar, N., M. Jay and R. Chemli. 2001. Chemical diversification trends in *Astragalus caprinus* (Leguminosae), based on the flavonoid pathway. Biochem. Syst. Ecol. 29(7): 727-738. Maps, Chemotaxonomy.
- Shukla, B.K. 2000. Studies on the pollination sub-systems in the five tree species of Caesalpiniaceae. In Mukerji, K.G., Chamola, B.P. and A.K. Sharma (eds.), Glimpses in botany: Professor G.S. Paliwal festschrift volume. New Delhi: APH, 2000 pp.18-28.
- Silva, V.N.T., de-Oliveira, F.M. and L.M. Conserva. 2001. Phenolic derivatives and terpenes from *Acosmium bijugum*. Biochem. Syst. Ecol. 29(11): 1189-1192. Chemotaxonomy.
- Simpson, B.B. 2000. *Hoffmannseggia humilis*, the correct name for *H. gladiata* (Fabaceae). Lundellia no.3: 41-42.
- Singh, V. 2001. Monograph on Indian subtribe *Cassiinae* (Caesalpiniaceae). Jodhpur: Scientific, 2001 xviii, 278pp. Icones, Maps, Chromosome numbers, Anatomy and Morphology, Keys.
- Sobrinho, E., A. Hervella, J.L. Ceresuela, A. Barbado, A. Viviani, F. de Andres and J.L. Tenorio. 2000. Morfologia y taxonomia de la seccion Dendrotelis del genero *Medicago* (Fabaceae). Portugal. Acta Biol. 19(1-4): 225-237. Chromosome numbers, Anatomy and Morphology.
- Soepadmo, E., L.G. Saw (eds.). 2000. Tree flora of Sabah and Sarawak: volume three. Kuala Lumpur: Forest Research Institute Malaysia, Sabah Forestry Department and Sarawak Forestry Department, 2000 xvi, 511pp. Icones, Anatomy and Morphology, Keys.
- SOKOLOFF, D.D. 2001. New records of *Lotus* (Leguminosae: Papilionoideae: Loteae) from Africa and southwest Asia. Kew Bull. 56(3): 751-720. Anatomy and Morphology, Keys.

- Sousa-Costa, N.M. and C. van den Berg. 2001. A new species of *Stylosanthes* Sw. (Leguminosae-Papilionoideae) from Guerrero, Mexico. Kew Bull. 56(1): 239-241. Icones, Anatomy and Morphology, Molecular systematics. *Stylosanthes salina* sp. nov.
- SOUSA, M.S. 2000. Especie nueva de *Lonchocarpus* (Leguminosae) de Nicaragua y Costa Rica. An. Inst. Biol. Univ. Nac. Auton. Mex., Bot. 70(2): 137-140. Icones, Anatomy and Morphology. *Lonchocarpus chiangii* sp. nov.
- SOUSA M.S., M. Ricker and H.M. Hernandez. 2001. Tree species of the family Leguminosae in Mexico. Harvard Pap. Bot. 6(1): 339-365.
- SPRENT, J.I. 2001. Nodulation as a taxonomic tool. HERENDEEN P.S. and A. BRUNEAU (eds.), Advances in legume systematics: part 9. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.21-43. Icones, Anatomy and Morphology.
- Srisanga, P. and S. Sasirat. 2000. *Acacia tonkinensis* I.C. Nielsen (Leguminosae-Mimosoideae), a new record for Thailand. Thai Forest Bull., Bot. no.28: 25-31. Icones, Maps, Anatomy and Morphology, Keys.
- Sri-Rama-Murthy, S., S. Sandhya-Rani and T. Pullaiah. 2000. A contribution to the flora of Eastern Ghats, India genus *Rhynchosia* Lour. (Faboideae-Fabaceae). In Gupta BK, ed. Higher plants of Indian subcontinent, vol. IX. Dehra Dun: Bishen Singh Mahendra Pal Singh, 2000 pp.1-15. (Indian Journal of Forestry Additional series; 11.) Icones, Maps, Keys.
- Stiefkens, L.B., G. Bernardello and G.J. Anderson. 2001. The somatic chromosomes of *Sophora fernandeziana* (Fabaceae), an endemic tree from Robinson Crusoe Island. Pac. Sci. 55(1): 71-75. Chromosome numbers.
- SYTIN, A.K. 2001. Generis *Astragalus* L. (Fabaceae) species nova e Russia Europaea.) Novosti Sist. Vyssh. Rast. 33: 132-134. Icones, Anatomy and Morphology. *Astragalus jelenevskyi* sp. nov.
- Tabur, S., S. Civelek and E. Bagci. 2001. Cytotaxonomic studies on some *Vicia* L. species growing in eastern Mediterranean and southern Aegean regions, I. Acta Bot. Gallica 148(2): 159-174. Maps, Chromosome numbers, Anatomy and Morphology.
- Tandon, R., K.R. Shivanna and H.Y. Mohan-Ram. 2001. Pollination biology and breeding system of *Acacia senegal*. Bot. J. Linn. Soc. 135(3): 251-262. Icones, Maps, Anatomy and Morphology, Reproductive biology.
- Tang, Y.P., J.H. Wang, Y.F. Li, F.C. Lou and W. Ma. 2001. (Flavonols and flavonol glycosides from the pericarp of *Sophora japonica* L.) J. Pl. Resour. Environ. 10(2): 59-60. Chemotaxonomy.
- Tapia-Pastrana, F. and P. Mercado-Ruaro. 2001. A combination of the "squash" and "splash" techniques to obtain the karyotype and assess meiotic behaviour of *Prosopis laevigata* L. (Fabaceae: Mimosoideae). Cytologia (Japan) 66(1): 11-17. Chromosome numbers.
- Teixeira-S-de, P. and A.C. Gabrielli. 2000. Anatomia de eixo vegetativo de *Dahlstedtia pinnata* (Benth.) Malme e *D. pentaphylla* (Taub.) Burk. (Leguminosae, Papilionoideae). (Vegetative axis anatomy of *Dahlstedtia pinnata* (Benth.) Malme and *D. pentaphylla* (Taub.) Burk. (Leguminosae, Papilionoideae).) Rev. Brasil. Bot. 23(1): 1-11. Icones, Anatomy and Morphology.
- THOMPSON, I.R. 2001. Morphometric analysis and revision of eastern Australian *Hovea* (Brongniartieae-Fabaceae). Austral. Syst. Bot. 14(1): 1-99. Icones, Maps, Anatomy and Morphology, Keys. Many new taxa.
- THOMPSON, I.R., P.Y. Ladiges and J.H. ROSS. 2001. Phylogenetic studies of the tribe Brongniartieae (Fabaceae) using nuclear DNA (ITS-1) and morphological data. Syst. Bot. 26(3): 557-570. Maps, Molecular systematics.
- Thseng, F.S., T.K. Lin and S.T. Wu. 2000. The relations of genus *Glycine* subgenus *Soja* and *Glycine formosana* Hosok. collected from Taiwan: revealed by RAPD analysis. J. Jap. Bot. 75(5): 270-279. Chromosome numbers, Molecular systematics.
- Thulin, M. and A.N. Al-Gifri. 2000. Two new species of *Acacia* (Leguminosae-Mimosoideae) from Yemen. Nordic J. Bot. 20(6): 691-696. Icones, Anatomy and Morphology.
- Thulin, M. and M. LAVIN. 2001. Phylogeny and biogeography of the *Ormocarpum* group (Fabaceae): a new genus *Zygocarpum* from the Horn of Africa region. Syst. Bot. 26(2): 299-317. Icones, Maps, Anatomy and Morphology, Keys, Molecular systematics. Many comb. nov.
- Torres-Colin, R. 2001. Una especie nueva de *Bauhinia* (Fabaceae: Caesalpinioideae: Cercideae) de Panama. Novon 11(1): 143-145. Icones, Anatomy and Morphology. *Bauhinia coclensis* sp. nov.

- Torres-Gonzalez, A.M., O. Toro-Chica and D.G. DEBOUCK. 2001. *Phaseolus talamancensis*, a new wild bean species (Leguminosae, Phaseolinae) from montane forests of eastern Costa Rica. *Novon* 11(2): 280-286. Icones, Maps, Anatomy and Morphology, Palynology.
- Trannin, I.C.B.; MOREIRA, F.M.S. and J.O. Siqueira. 2001. Growth and nodulation of *Acacia mangium*, *Enterolobium contortsiliquum* and *Sesbania virgata* in heavy metal contaminated soil. *Revista Brasileira de Ciência do Solo*, 25: 743-753.
- Trannin, I.C.B.; MOREIRA, F.M.S.; Siqueira, J.O. and A. Lima. 2001. Tolerance of *Bradyrhizobium* e de *Azorhizobium* strains to copper, cadmium and zinc "in vitro". *Revista Brasileira de Ciência do Solo*, 25: 305-316.
- TUCKER, S.C. 2001. Organ loss in detarioid and other leguminous flowers, and the possibility of saltatory evolution. In HERENDEEN, P.S. and A. BRUNEAU (eds.), *Advances in legume systematics: part 9*. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.107-120. Icones, Anatomy and Morphology.
- TURNER, B.L. 2000. The Texas species of *Calliandra* (Leguminosae, Mimosoideae). *Lundellia* no.3: 13-18. Icones, Maps, Anatomy and Morphology, Keys. *Calliandra iselyi* sp. nov.
- ULIBARRI, E.A. 2000. Una nueva especie de *Adesmia* (Leguminosae-Papilionoideae) de Bolivia. (A new species of *Adesmia* (Leguminosae-Papilionoideae) from Bolivia.) *Hickenia* 3(16-23): 57-59. Icones, Anatomy and Morphology. *Adesmia kieslingii* sp. nov.
- ULIBARRI, E.A. and A. Burkart. 2000. Sinopsis de las especies de *Adesmia* (Leguminosae, Adesmieae) de la Argentina. (Synopsis of the species of *Adesmia* (Leguminosae, Papilionoideae) of Argentina.) *Darwiniana* 38(1-2): 59-126. Icones, Chromosome numbers, Anatomy and Morphology, Keys. 2 var. nov.; 3 comb. nov.
- Ulziikhutag, N. 2001. (Sectio nova et species novae subgeneris *Cercidothrix* Bunge generis *Astragalus* L. (Fabaceae) ex Asia Centrali. *Novosti Sist. Vyssh. Rast.* 33: 135-145. Anatomy and Morphology. 6 spp. nov.
- Valdes-Castrillon, B. 2000. Sobre la presencia de *Hedysarum hedysaroides* (L.) Schinz et Thell. (Leguminosae) en la Peninsula Iberica. (On the presence of *Hedysarum hedysaroides* (L.) Schinz and Thell. (Leguminosae) from the Iberian Peninsula.) *An. Jard. Bot. Madrid* 57(2): 453-454.
- Vela, E., R. Giraud, Y. Morvan and F. Medail. 2000. A propos d' *Astragalus tragacantha* L. [syn.: *A. massiliensis* (Miller) Lam.], espece protegee: stations nouvelles sur la Commune de Marseille. *Monde Pl.* 95(470): 10-11.
- Verdcourt, B. 2001. A new species of *Vigna* (Leguminosae-Phaseoleae) from Bolivia. *Kew Bull.* 56(1): 229-232.
- Vivant, J. 2000. *Medicago orbicularis* subsp. *castellana* (J. Casellas) P. Monts. observe dans les Hautes-Pyrenees en Vallee d'Aure. *Monde Pl.* 95(470): 8.
- Wang, Q. and C.Y. Yang. 2001. (A new species of the genus *Oxytropis* from Xinjiang, China.) *Acta Phytotax. Sin.* 39(2): 151-153. Icones, Anatomy and Morphology. *Oxytropis pseudohirsuta* sp. nov.
- Wang, Q., C.Y. Yang, X.Y. Zhu and H. OHASHI. 2001. A new species of *Oxytropis* DC. (Leguminosae) from Xinjiang, China. *J. Jap. Bot.* 76(4): 219-221. Icones, Anatomy and Morphology, Keys. *Oxytropis fragiliphylla* sp. nov.
- Watson, L.E., H. Sayed-Ahmed and A. Badr. 2000. Molecular phylogeny of Old World *Trifolium* (Fabaceae), based on plastid and nuclear markers. *Pl. Syst. Evol.* 224(3-4): 153-171. Molecular systematics.
- Welsh, S.L. and N.D. Atwood. 2001. New taxa and nomenclatural proposals in miscellaneous families: Utah and Arizona. *Rhodora* 103(913): 71-95. Icones, Chromosome numbers, Anatomy and Morphology, Keys. Many new taxa.
- Wilson, P.G. 2001. Some name changes in *Indigofera* (Fabaceae). *Taxon* 50(2): 491-493. 3 comb. nov.
- WOJCIECHOWSKI, M., M.J. SANDERSON, K.P. Steele and A. LISTON. 2001. Molecular phylogeny of the "temperate herbaceous tribes" of papilionoid legumes: a supertree approach. In Herendeen P.S. and A. Bruneau (eds.), *Advances in legume systematics: part 9*. Kew: Royal Botanic Gardens, Kew, 2000 publ. 2001 pp.277-298. Molecular systematics.
- Zamora, N. 2000. Nuevas especies y combinaciones en leguminosas de Mesoamerica. *Novon* 10(2): 175-180. Icones, Anatomy and Morphology, Keys. 2 spp. nov.; 2 comb. nov.

Zanin, L.A. and M.A. Cangiano. 2001. El cariotipo de *Hoffmannseggia glauca* (Fabaceae). (The karyotype of *Hoffmannseggia glauca* (Fabaceae).) *Darwiniana* 39(1-2): 11-13. Chromosome numbers.

Zarre-Mobarakeh, S. 2000. Systematic revision of *Astragalus* sect. *Adiaspastus*, sect. *Macrophyllum* and sect. *Pterophorus* (Fabaceae). *Englera* 18: 219pp. Icones, Maps, Chromosome numbers, Anatomy and Morphology, Keys.

ZARRE, S. and D. PODLECH. 2001. Taxonomic revision of *Astragalus* sect. *Acanthophaea* (Fabaceae). *Sendtnera* 7: 233-251. Icones, Anatomy and Morphology, Keys. 2 spp. nov.

ZHANG, M.L. 2000. Studies on geographical distribution pattern of the subgenus *Pogonophaea* (Fabaceae: *Astragalus*) in China using GIS technique. *Acta Bot. Sin.* 42(8): 849-854. Maps.

Zhao, D.L., Z. HU, W. Chen, X.W. Jiang and L.S. Wang. 2001. Karyotype analysis of *Sphallerocarpus gracilis* and *Trifolium pratense* from China. *Acta Bot. Bor. Occid. Sin.* 21(5): 1026-1030. Chromosome numbers.

Zhu, C.S., W.X. Su and Z.L. Yuan. 2000. A revision of *Lathyrus* and *Campylotropis* (Leguminosae) in Henan.) *J. Wuhan Bot. Res.* 18(6): 471-472.

Zhu, X.Y., Y.F. Du and H. OHASHI. 2000. A new *Oxytropis* species (Leguminosae) from Xinjiang in China. *J. Jap. Bot.* 75(5): 289-291. Icones, Anatomy and Morphology, Keys. *Oxytropis tashkurensis* sp. nov.

Zhu, X.Y. and H. OHASHI. 2001. *Dendrolobium rugosum* (Prain) Schindler (Leguminosae), a new record to China. *Acta Phytotax. Sin.* 39(5): 477-480. Icones, Anatomy and Morphology.

Zhu, Y.Q., A.P. Zhou, Y.S. Li, B. Li and J.F. Xu. 2000. New species from Shandong. *Bull. Bot. Res. (China)* 20(1): 7-8. Icones, Anatomy and Morphology. *Vicia villosa* var. *alba* var. nov.